

Fishery Data Series No. 07-39

Migratory Timing And Abundance Estimates Of Sockeye Salmon Into Upper Cook Inlet, Alaska, 2005

by

Pat Shields

and

Mark Willette

June 2007

Alaska Department of Fish and Game

Division of Sport Fish and Commercial Fisheries



Symbols and Abbreviations

The following symbols and abbreviations, and others approved for the Système International d'Unités (SI), are used in Division of Sport Fish Fishery Manuscripts, Fishery Data Series Reports, Fishery Management Reports, and Special Publications without definition. All others must be defined in the text at first mention, as well as in the titles or footnotes of tables and in figures or figure captions.

| | | | | | |
|---------------------------------------|--------------------|--------------------------|----------------------------------|----------------------------------|-------------------------|
| Weights and measures (metric) | | General | | Measures (fisheries) | |
| centimeter | cm | Alaska Administrative | | fork length | FL |
| deciliter | dL | Code | AAC | mid-eye-to-fork | MEF |
| gram | g | all commonly accepted | | mid-eye-to-tail-fork | METF |
| hectare | ha | abbreviations | e.g., Mr., Mrs., AM, PM, etc. | standard length | SL |
| kilogram | kg | | | total length | TL |
| kilometer | km | all commonly accepted | | | |
| liter | L | professional titles | e.g., Dr., Ph.D., R.N., etc. | Mathematics, statistics | |
| meter | m | | | <i>all standard mathematical</i> | |
| milliliter | mL | at | @ | <i>signs, symbols and</i> | |
| millimeter | mm | compass directions: | | <i>abbreviations</i> | |
| | | east | E | alternate hypothesis | H _A |
| Weights and measures (English) | | north | N | base of natural logarithm | <i>e</i> |
| cubic feet per second | ft ³ /s | south | S | catch per unit effort | CPUE |
| foot | ft | west | W | coefficient of variation | CV |
| gallon | gal | copyright | © | common test statistics | (F, t, χ^2 , etc.) |
| inch | in | corporate suffixes: | | confidence interval | CI |
| mile | mi | Company | Co. | correlation coefficient | |
| nautical mile | nmi | Corporation | Corp. | (multiple) | R |
| ounce | oz | Incorporated | Inc. | correlation coefficient | |
| pound | lb | Limited | Ltd. | (simple) | r |
| quart | qt | District of Columbia | D.C. | covariance | cov |
| yard | yd | et alii (and others) | et al. | degree (angular) | ° |
| | | et cetera (and so forth) | etc. | degrees of freedom | df |
| Time and temperature | | exempli gratia | | expected value | <i>E</i> |
| day | d | (for example) | e.g. | greater than | > |
| degrees Celsius | °C | Federal Information | | greater than or equal to | ≥ |
| degrees Fahrenheit | °F | Code | FIC | harvest per unit effort | HPUE |
| degrees kelvin | K | id est (that is) | i.e. | less than | < |
| hour | h | latitude or longitude | lat. or long. | less than or equal to | ≤ |
| minute | min | monetary symbols | | logarithm (natural) | ln |
| second | s | (U.S.) | \$, ¢ | logarithm (base 10) | log |
| | | months (tables and | | logarithm (specify base) | log ₂ , etc. |
| Physics and chemistry | | figures): first three | | minute (angular) | ' |
| all atomic symbols | | letters | Jan, ..., Dec | not significant | NS |
| alternating current | AC | registered trademark | ® | null hypothesis | H ₀ |
| ampere | A | trademark | ™ | percent | % |
| calorie | cal | United States | | probability | P |
| direct current | DC | (adjective) | U.S. | probability of a type I error | |
| hertz | Hz | United States of | | (rejection of the null | |
| horsepower | hp | America (noun) | USA | hypothesis when true) | α |
| hydrogen ion activity | pH | U.S.C. | United States | probability of a type II error | |
| (negative log of) | | | Code | (acceptance of the null | |
| parts per million | ppm | U.S. state | use two-letter | hypothesis when false) | β |
| parts per thousand | ppt, ‰ | | abbreviations | second (angular) | " |
| | | | (e.g., AK, WA) | standard deviation | SD |
| volts | V | | | standard error | SE |
| watts | W | | | variance | |
| | | | | population | Var |
| | | | | sample | var |

FISHERY DATA SERIES NO. 07-39

**MIGRATORY TIMING AND ABUNDANCE ESTIMATES OF SOCKEYE
SALMON INTO UPPER COOK INLET, ALASKA, 2005**

by

Pat Shields and Mark Willette

Alaska Department of Fish and Game, Division of Commercial Fisheries, Soldotna

Alaska Department of Fish and Game
Division of Sport Fish, Research and Technical Services
333 Raspberry Road, Anchorage, Alaska, 99518

June 2007

The Division of Sport Fish Fishery Data Series was established in 1987 for the publication of technically oriented results for a single project or group of closely related projects. Since 2004, the Division of Commercial Fisheries has also used the Fishery Data Series. Fishery Data Series reports are intended for fishery and other technical professionals. Fishery Data Series reports are available through the Alaska State Library and on the Internet: <http://www.sf.adfg.state.ak.us/statewide/divreports/html/intersearch.cfm> This publication has undergone editorial and peer review.

Pat Shields and Mark Willette

*Alaska Department of Fish and Game, Division of Commercial Fisheries,
43961 Kalifornsky Beach Road, Suite B, Soldotna, AK 99669-8367, USA*

This document should be cited as:

Shields, P., and M. Willette. 2007. Migratory timing and abundance estimates of sockeye salmon into Upper Cook Inlet, Alaska, 2005. Alaska Department of Fish and Game, Fishery Data Series No. 07-39, Anchorage.

The Alaska Department of Fish and Game (ADF&G) administers all programs and activities free from discrimination based on race, color, national origin, age, sex, religion, marital status, pregnancy, parenthood, or disability. The department administers all programs and activities in compliance with Title VI of the Civil Rights Act of 1964, Section 504 of the Rehabilitation Act of 1973, Title II of the Americans with Disabilities Act (ADA) of 1990, the Age Discrimination Act of 1975, and Title IX of the Education Amendments of 1972.

If you believe you have been discriminated against in any program, activity, or facility please write:

ADF&G ADA Coordinator, P.O. Box 115526, Juneau AK 99811-5526

U.S. Fish and Wildlife Service, 4040 N. Fairfax Drive, Suite 300 Webb, Arlington VA 22203

Office of Equal Opportunity, U.S. Department of the Interior, Washington DC 20240

The department's ADA Coordinator can be reached via phone at the following numbers:

(VOICE) 907-465-6077, (Statewide Telecommunication Device for the Deaf) 1-800-478-3648, (Juneau TDD) 907-465-3646, or (FAX) 907-465-6078

For information on alternative formats and questions on this publication, please contact:

ADF&G, Sport Fish Division, Research and Technical Services, 333 Raspberry Road, Anchorage AK 99518 (907)267-2375.

TABLE OF CONTENTS

| | Page |
|---|-------------|
| LIST OF TABLES..... | ii |
| LIST OF FIGURES..... | ii |
| LIST OF APPENDICES..... | ii |
| ABSTRACT..... | 1 |
| INTRODUCTION..... | 1 |
| METHODS..... | 1 |
| Test Fishing..... | 1 |
| Describing the Salmon Migration and Projecting Total Run..... | 3 |
| RESULTS AND DISCUSSION..... | 6 |
| ACKNOWLEDGEMENTS..... | 10 |
| REFERENCES CITED..... | 11 |
| TABLES AND FIGURES..... | 15 |
| APPENDIX A..... | 29 |

LIST OF TABLES

| Table | Page |
|--|-------------|
| 1. Summary of sockeye salmon fishing effort, daily and cumulative catch, and daily and cumulative CPUE, Upper Cook Inlet offshore test fish project, 2005. | 16 |
| 2. Estimated sockeye salmon catch by date and station, Upper Cook Inlet offshore test fish project, 2005. | 17 |
| 3. Estimated sockeye salmon CPUE by date and station, Upper Cook Inlet offshore test fish project, 2005. | 18 |
| 4. Total run estimates for sockeye salmon to Upper Cook Inlet, Alaska, made during the 2005 season. | 19 |
| 5. Estimated total Kenai River sockeye run (millions) in 2005 estimated from total offshore test fish CPUE and age composition stock allocation estimates through July 21, and July 27. | 21 |
| 6. A comparison of methods used to make post-season adjustments to the offshore test fish final CPUE. | 22 |
| 7. Mean date of the sockeye salmon run across the Anchor Point transect, Upper Cook Inlet offshore test fish project, 1979–2005. | 23 |
| 8. Sockeye salmon age-composition, mean length (mm), and number of samples by station, Upper Cook Inlet offshore test fish project, 2005. | 24 |

LIST OF FIGURES

| Figure | Page |
|---|-------------|
| 1. Location of fishing districts and offshore test fish transect in Cook Inlet, Alaska 2005. | 25 |
| 2. Linear regression of the relationship between offshore test fish unadjusted cumulative CPUE and Upper Cook Inlet sockeye salmon total annual run. | 26 |
| 3. Cumulative proportions estimated for the sockeye salmon run to Upper Cook Inlet, Alaska, 2005. | 27 |
| 4. OTF error in forecasting the total run of sockeye salmon to Upper Cook Inlet using the 20-July best-fit estimate. | 28 |

LIST OF APPENDICES

| Appendix | Page |
|---|-------------|
| A1. Summary of pink salmon fishing effort, daily and cumulative catch, and daily and cumulative CPUE, Upper Cook Inlet offshore test fish project, 2005. | 30 |
| A2. Estimated pink salmon catch by date and station, Upper Cook Inlet offshore test fish project, 2005. | 31 |
| A3. Estimated pink salmon CPUE by date and station, Upper Cook Inlet offshore test fish project, 2005. | 32 |
| A4. Summary of chum salmon fishing effort, daily and cumulative catch, and daily and cumulative CPUE, Upper Cook Inlet offshore test fish project, 2005. | 33 |
| A5. Estimated chum salmon catch by date and station, Upper Cook Inlet offshore test fish project, 2005. | 34 |
| A6. Estimated chum salmon CPUE by date and station, Upper Cook Inlet offshore test fish project, 2005. | 35 |
| A7. Summary of coho salmon fishing effort, daily and cumulative catch, and daily and cumulative CPUE, Upper Cook Inlet offshore test fish project, 2005. | 36 |
| A8. Estimated coho salmon catch by date and station, Upper Cook Inlet offshore test fish project, 2005. | 37 |
| A9. Estimated coho salmon CPUE by date and station, Upper Cook Inlet offshore test fish project, 2005. | 38 |
| A10. Summary of Chinook salmon fishing effort, daily and cumulative catch, and daily and cumulative CPUE, Upper Cook Inlet offshore test fish project, 2005. | 39 |
| A11. Estimated Chinook salmon catch by date and station, Upper Cook Inlet offshore test fish project, 2005. | 40 |
| A12. Estimated Chinook salmon CPUE by date and station, Upper Cook Inlet offshore test fish project, 2005. | 41 |
| A13. Entry pattern of sockeye salmon into Upper Cook Inlet, Alaska, 2005 estimated from daily CPUE measured at the latitude of Anchor Point. | 42 |
| A14. Chemical and physical observations made in Upper Cook Inlet, Alaska, during the 2005 offshore test fish project. | 43 |
| A15. Yearly mean values of physical observations made during the 1995–2005 offshore test fish project. | 48 |
| A16. Yearly mean values for selected chemical and physical variables collected during the offshore test fish project, 1979–2005. | 50 |

ABSTRACT

A test fishery was conducted during the 2005 Upper Cook Inlet (UCI) commercial salmon fishery. The primary objective of the test fishery was to estimate the abundance and timing of the sockeye salmon *Oncorhynchus nerka* run, as measured along a transect near the southern boundary of the UCI management area. The test fishery was conducted from 1 July to 30 July and captured 4,108 sockeye salmon, representing 2,643 catch per unit of effort (CPUE) points. The midpoint of the 2005 run occurred on 21 July, which was 6 days late relative to the historical mean date of 15 July. This represents the latest run-timing since the test fishery began in 1979. A non-linear mathematical model estimated the 2005 test fish project spanned approximately 68.8% of the sockeye salmon run. The test fish final passage rate was estimated at approximately 2,500 sockeye salmon per CPUE point. Two formal estimates of the size and timing of the 2005 sockeye salmon run were made during the commercial fishing season, with the first best-fit estimator from each analysis forecasting a total run to UCI of 9.2 and 8.2 million sockeye salmon, respectively. These estimates deviated from the preliminary total run estimate of 7.8 million by 18% and 5%, respectively. The test fish project once again provided valuable data used to aid in critical inseason commercial fishery management decisions.

Key words: Salmon, *Oncorhynchus*, Upper Cook Inlet, Alaska, test fishery, migratory behavior.

INTRODUCTION

In 1979 the Alaska Department of Fish and Game (ADF&G) began an Offshore Test Fish (OTF) project near the southern boundary of the Upper Cook Inlet (UCI) salmon management area (Figure 1). The objective of the project has been to estimate the total run and run-timing of sockeye salmon *Oncorhynchus nerka* returning to UCI during the commercial salmon fishing season. These data have become extremely important to ADF&G management biologists as they help set and adjust commercial fishing times and areas to most efficiently harvest sockeye salmon that are surplus to spawning needs. Moreover, the Alaska Board of Fisheries (BOF) has assembled management plans which require inseason estimates of the size of the sockeye salmon run in order to implement specific components of the various plans. The OTF project has increasingly become one of the most important tools fishery managers utilize to make inseason fishery management decisions.

Test fishing results have been reported annually since 1979 (Hilsinger 1988; Hilsinger and Waltemyer 1987; Shields 2000, 2001, 2003; Shields and Willette 2004, 2005; Tarbox 1990-1991, 1994-1999; Tarbox and King 1992; Tarbox and Waltemyer 1989; Waltemyer 1983a, b, 1986a, b). This report presents the results of the 2005 test-fishing project.

METHODS

TEST FISHING

Sockeye salmon returning to UCI were sampled by fishing 6 geographically fixed stations between Anchor Point and the Red River Delta (Figure 1). Stations were numbered consecutively from east to west, with station locations (latitude and longitude) determined with global positioning system technology. A chartered test-fishing vessel, *FV Americanus*, sampled all 6 stations (numbered 4, 5, 6, 6.5, 7, and 8) daily, traveling east to west on odd-numbered days and west to east on even-numbered days. Sampling started on 1 July and continued through 30 July. The vessel fished 366 m (1,200 ft or 200 fathoms) of 13 cm (5 1/8 in) multi-filament drift

gillnet. The net was 45 meshes deep and constructed of double knot Super Crystal shade number 1, with a filament size of number 53/S6F¹.

The following physical and chemical readings were taken at the start of each set: air temperature, water temperature (at 1 m below the surface), wind velocity and direction, tide stage, water depth, and water clarity. Air and water temperatures were measured using a YSI salinity/temperature meter. Wind speed was measured in knots and direction was recorded as 0 (no wind), 1 (north), 2 (northeast), 3 (east), 4 (southeast), 5 (south), 6 (southwest), 7 (west), or 8 (northwest). Tide stage was classified as 1 (high slack), 2 (low slack), 3 (flooding), or 4 (ebbing) by observing the movement of the vessel while drifting with the gill net. Water depth was measured in fathoms (fm) using a Simrad echo sounder, and water clarity was measured in meters (m) using a 17.5 cm secchi disk.

All salmon captured in the drift gillnet were enumerated and identified to species and sex. Sockeye salmon ($n \leq 30$ at each station) were measured for fork length (mideye to tail fork) to the nearest mm and also had a scale removed (for age determination) as described by Koo (1955). Scales were mounted on gum cards and impressions made in cellulose acetate, as described by Clutter and Whitesel (1956). The age of each fish was determined after examining scales with a microfiche viewer under 40x magnification. Ages were reported in European notation (Koo 1962) and followed criteria established by Mosher (1969) and Tobias et al. (1994).

The number of fish caught at each station (s) on each day (i) was expressed as a catch per unit of effort (CPUE) statistic, or index, and standardized to the number of fish caught in 100 fathoms of gear in 1 hour of fishing time.

$$CPUE_{s,i} = \frac{100 \text{ fm} \times 60 \text{ min} \times \text{number of fish}}{\text{fm of gear} \times MFT} \quad (1)$$

where:

MFT = mean fishing time.

Mean fishing time (MFT) was calculated as:

$$MFT = (C - B) + \frac{(B - A) + (D - C)}{2} \quad (2)$$

where:

A = time net deployment started,

B = time net fully deployed,

C = time net retrieval started, and

D = time net fully retrieved.

Once deployed at a station, the drift gillnet was fished 30 minutes before retrieval started. However, the net was capable of capturing fish prior to being fully deployed, and during the time it was being retrieved. MFT was therefore adjusted by summing the total time it took to set and retrieve the net,

¹ Product names used in this report are included for scientific completeness, but do not constitute a product endorsement.

then dividing this time in half, and adding it to the time when the entire net was deployed and fished.

Daily $CPUE_i$ data were summed for all m stations (typically 6) as follows:

$$CPUE_i = \sum_{s=1}^m CPUE_{s,i} \quad (3)$$

Cumulative $CPUE_i$ ($CCPUE_d$) was given by:

$$CCPUE_d = \sum_{i=1}^d CPUE_i \quad (4)$$

where:

d = day for which estimate is being made.

DESCRIBING THE SALMON MIGRATION AND PROJECTING TOTAL RUN

The run entry pattern was described for each of the previous years based on the respective test fishing data as per Mundy (1979):

$$Y_{yr,d} = 1 / (1 + e^{-(a+bd)}) \quad (5)$$

where:

$Y_{yr,d}$ = modeled cumulative proportion of $CCPUE_{yr,f}$ (f = final day of season) for year yr as of day d ,

a and b = model parameters.

Variables without the subscript yr indicating year refer to the current year's estimate. To determine which of the previous run-timing models most closely fit the current year's data and to estimate total run for the entire season (TR_f), a projection of the current year's $CCPUE_d$ at the end of the season ($CCPUE_f$) was estimated as per Waltemyer (1983a):

$$CCPUE_f = \frac{\sum_{i=0}^n CCPUE_d^2}{\sum_{i=0}^n Y_{yr,d} \cdot CCPUE_d} \quad (6)$$

This model assumes that the average day of return and its variance for previous year yr is the same for the current season (Mundy 1979). To test this assumption, in-season Y_d was estimated as:

$$Y_d = \frac{CCPUE_d}{CCPUE_f} \quad (7)$$

and mean squared error (MSE) between Y_d and $Y_{yr,d}$ was estimated as:

$$MSE = \frac{\sum_{i=0}^n (Y_{yr,d} - Y_d)^2}{n+1} \quad (8)$$

where:

$n+1$ = number of days in the comparison.

Years were ranked from lowest MSE (best model) to highest (worst), and the best fit years were used to estimate $CCPUE_f$. Catchability, or the fraction of the available population taken by a defined unit of fishing effort, was estimated as:

$$q_d = \frac{CCPUE_d}{r_d} \quad (9)$$

where:

q_d = estimated cumulative catchability as of day d ,

r_d = cumulative total run as of day d .

The cumulative total run on day d was estimated from the sum of all commercial, recreational, and personal use harvests to date, the estimated total escapement to date, and an estimate of the number of sockeye salmon residual in the district at that time. Commercial harvest data is estimated inseason from catch reports called or faxed into the ADF&G office. All commercially harvested salmon in UCI, whether sold or kept for personal use, are required to be reported to the Soldotna ADF&G office by fishermen or the processors they sell their fish to within 12 hours of the close of a fishing period. For a complete list of reporting requirements, please see the following statute: 16.05.690(a) and regulation: 5 AAC 39.130. Recreational harvest data was estimated inseason and provided by Division of Sport Fish staff. Personal use harvests were also estimated inseason from daily reports from the various fisheries in combination with an assessment of previous year's personal use catches from runs of similar abundance. Total escapement to date included estimated escapements into all monitored systems (Crescent, Susitna, Kenai and Kasilof Rivers, and Fish Creek) and unmonitored systems, which are assumed to be 15% of the escapement into monitored systems (Tobias and Willette 2003). The number of fish residual in the district was estimated by assuming exploitation rates of 70% in set net fisheries, 40% in district-wide drift net fisheries, and 25% in reduced district drift net fisheries (Mundy et al. 1993). For example, if the drift gillnet fleet harvested 500 thousand sockeye salmon on an inlet wide fishing period, the number of sockeye salmon originally in the district would have been estimated at 1,250 thousand ($500/0.40 = 1,250$) and the number remaining, or the residual, would have been estimated at 750 thousand ($1,250 - 500 = 750$).

Passage rate, the expansion factor used to convert CPUE into estimated numbers of salmon passing the test fishing transect, was calculated as:

$$PR_d = 1/q_d \quad (10)$$

Total run at the end of the season (TR_f) was then estimated from:

$$TR = PR \cdot CCPUE_f \quad (11)$$

To calculate the midpoint of the run, which is the day on which approximately 50% of the total run had passed the OTF transect, the following formula was used:

$$M = a/b \quad (12)$$

where:

M = mean date of run,

a and b = model parameters.

Because the test fishery did not encompass the entire sockeye salmon run, the total $CCPUE_f$ for the test fishery was estimated after the season using the following two methods:

$$CCPUE_f^h = CCPUE_f \cdot \frac{H_t}{H_L} \quad (13)$$

where:

$CCPUE_f^h$ = total estimated $CCPUE_f$ for the season, based on harvest,

H_t = total commercial harvest for the season,

H_L = total commercial harvest through final day of test fishery (f+2), and

L = number of days (lag time) it took salmon to travel from test fishery to commercial harvest areas (2 days).

$$CCPUE_t^r = CCPUE_f \cdot \frac{E_t + H_t}{\sum_{s=1}^6 E_L + H_L} \quad (14)$$

where:

$CCPUE_t^r$ = total estimated $CCPUE_f$ for the season, based upon total run,

E_t = total escapement for the season

H_t = total commercial harvest for the season

E_L = total Upper Cook Inlet escapement through final day of test fishery

H_L = total Upper Cook Inlet commercial harvest through final day of test fishery

L = number of days (lag time) it took salmon to travel from test fishery to spawning streams or to be available for commercial harvest.

The total run adjustment to $CCPUE_f$ (Equation 14) has replaced adjustments based on harvest alone (Equation 9) primarily due to modifications to commercial fishing management plans made by the BOF. Current management plans allow for much less fishing in August than in the past; therefore, adjustments based on harvest alone would not have accurately reflected the additional fish that entered the district after the test fishery ceased. The total run to date on the last day of the test fishery was computed by summing all commercial harvest data and estimates of escapement from the 4 sockeye salmon sonar enumeration sites, 1 weir site, and an estimate of escapement to all unmonitored systems through day d . An estimate of sockeye salmon

escapement to all non-monitored systems in UCI is considered to be 15% of the monitored runs. Lag times are the approximate time needed for fish to migrate from the test fish transect to a destination. As suggested by Mundy et al. (1993), lag times must be accounted for when estimating the total run passing the test fish transect on day d . A lag time of up to 2 days was assumed for fish harvested in the commercial fishery. The following lag times were assumed for fish entering the escapement: Crescent River, 1 day; Kasilof and Kenai rivers, 2 days; and Yentna River and Fish Creek, 7 days (15% of these totals are allocated to unmonitored systems) (Mundy et al. 1993). The number of sockeye salmon harvested in sport and personal use fisheries after test fishing ceased that are not already accounted for in escapement monitoring are assumed to be insignificant and therefore are not utilized in the $CCPUE_f$ post test fishery adjustment.

Adjusted estimates of $CCPUE_f$ ($CCPUE_t^h$ and $CCPUE_t^r$) were used for postseason estimates of TR_f .

RESULTS AND DISCUSSION

A total of 4,108 sockeye salmon were captured during the 2005 test fishery, as well as 186 pink salmon *O. gorbuscha*, 448 chum salmon *O. keta*, 546 coho salmon *O. kisutch*, and 8 Chinook salmon *O. tshawytscha* (Tables 1 and 2, Appendices A1–A12). The test boat was unable to fish 2 of the 6 stations on 10 July due to rough water, therefore catch data for these days were interpolated by averaging 9 July and 11 July catch data from these 2 stations. Sockeye salmon daily cumulative catches ranged from 25 to 467 fish (Table 1). The unadjusted total sockeye salmon $CCPUE_f$ for the 2005 project was 2,643, with daily CPUE values ranging from 19 to 208. The $CCPUE_f$ of 2,643 represented the highest final CPUE value ever measured in the OTF program, yet the total estimated run in 2005 was only the third strongest during this same time period. In 1992, the number of stations sampled by the test fish boat was standardized to the current 6 still being fished (Tarbox 1994). Since that time, the relationship between the annual test fish unadjusted $CCPUE_f$ and the total annual run of sockeye salmon to UCI has been significantly correlated ($P < 0.05$), but the fit of the data ($R^2 = 0.29$) is not strong enough to utilize test fish $CCPUE_f$ alone to predict the total annual run (Figure 2). Moreover, the autocorrelation and partial autocorrelation functions of the regression residuals indicated no significant ($p < 0.05$) correlations at lags up to 5 years. However, the annual $CCPUE_f$ statistic is reliable enough to provide a rough gauge of annual run strength, as the regression formula averaged about 27% error in predicting the size of the annual run from the true value.

The distribution of sockeye salmon catches along the test fish transect was similar to the distribution of CPUE values (Tables 2 and 3), which would be expected when fishing occurs at fixed intervals at each station.

Tarbox and Waltemeyer (1989) provided further detail into some of the assumptions the curve fitting procedures utilized to estimate the total CPUE statistic during the season. One of the major assumptions is that 24 June represents the first day of the sockeye salmon run to UCI. Variability in actual runs can therefore result in an average or early run being misclassified as late, especially during the first couple weeks of the test fish program. For this reason, 20 July has been the approximate date that commercial fishery staff have used for their first formal estimate of each year's total run size. By this time in the run, there are enough data points in the

current year's run-timing curve to provide a more accurate estimate of what the total CPUE will be at the end of the season. In addition, Tarbox and King (1992) and later OTF annual reports all suggested that the initial first best fit estimate made around mid July was not always the most accurate, i.e., the second or third best fit estimate should carefully be considered. Therefore, the method now used to make the first formal inseason estimate of the total run includes an examination of the top five or six best fits. Careful consideration is given to those years whose fits reveal the least day to day change in predicting what the final test fish CPUE will be. These years are identified as being potentially better fits, especially if the mean sum of squares statistic is also improving. Sockeye salmon run-timing from other areas of the state has also been considered to see if a consistent pattern exists. Until more years of data are collected from various runs, caution is advised when choosing the top fits to make projections of the final test fish CPUE, especially prior to 20 July.

The first formal estimate of the 2005 total run of sockeye salmon to UCI was made after the 21 July inlet-wide drift gillnet fishing period (Table 4). The 2005 test fish *CCPUE* curve was mathematically compared (mean sum of squares statistic) to runs from 1979 through 2004, with fits of the data ranked from best to worst. Based on an estimate of the run to date of 4.4 million sockeye salmon through 21 July (this estimate included an estimate of the number of fish residual in the district based upon assumed exploitation rates by different gear types), a passage rate of 3,009 was calculated. The best fit of the current year's *CCPUE* curve tracked the 1994 run, which was a 4-day late run, and estimated a total run of 9.17 million fish. As cautioned earlier, the first best fit using data from approximately 20 July often turns out not to be the best fit by year's end, so the top five best fits were considered. From these data, a total run estimate of 5.8 to 9.2 million was determined. The top three best fits all tracked runs that were 1 or more days late relative to the 15 July mean date of entry at the test fish transect line.

The next formal estimate of the total run of sockeye salmon to UCI in 2005 was made following the 25 July fishing period (Table 4). The run to date was now estimated at 5.3 million fish, with a *CCPUE* of 1,922. The passage rate was therefore estimated at 2,769 fish per CPUE point. The best fit still tracked the 1994 run, which projected a *CCPUE* of 2,947 and a total run of 8.2 million fish. The top five best fits now all tracked late runs and projected a total run to UCI of 5.9 to 8.2 million fish. The actual total run estimate of sockeye salmon to UCI in 2005 was 7.8 million fish, which includes commercial, sport, and personal use harvests, as well as escapement to all systems. Therefore, the first best fit run estimates from the two formal inseason forecasts of the 2005 run size were approximately 18% and 5% higher than the actual run size, respectively.

Figure 4 depicts the OTF error in projecting the total sockeye salmon run based on the best fits of the data on or soon after 20 July. As can be seen in this figure, the error in the 20 July estimate has been significant (>30%) only on runs that were 2 or more days early. In fact, for runs that are 1-day early, on time, or late, the error in the 20 July estimate of the total run ranges from -6.4% to +17.6% (average = +5.3%) of the actual run. Conversely, for runs that enter the district 2 or more days earlier than average, the OTF curve-fitting estimator does not perform nearly as well, with a range in error of +8.8% to +75.4%, or an average of +36.5%. It is quite likely that as additional data representing runs with more variable run timing are added to the database, the OTF projections will become more accurate for early runs.

UCI commercial fishery management plans compel ADF&G to make an inseason estimate of the number of sockeye salmon in each year's run that are of Kenai River origin. Various management

actions in both sport and commercial fisheries are tied to the total abundance of Kenai River sockeye salmon, which is characterized by three different ranges: less than 2 million fish, between 2 and 4 million fish, and greater than 4 million fish (Shields 2006). Based on commercial, sport, and personal use harvest, and escapement data through 21 July, the total sockeye salmon run through this date of Kenai River origin was estimated at 2.6 million fish (Table 5). Again, using the top five best fits of the current year's *CCPUE* curve to estimate the total sockeye salmon run, and assuming that 65% of the remaining run would be of Kenai River origin, the estimated total Kenai River run for 2005 was estimated at 3.7 to 5.9 million fish. The 25 July run projection from the top five best fits from the test fishery estimated that the total Kenai River sockeye salmon run would be 3.9 to 5.5 million fish. These formal run projections of the size of the Kenai River sockeye salmon run were critical to inseason commercial fishery management decisions, as management plans are tied to the size of the Kenai River sockeye salmon run. For example, if the size of the Kenai River run exceeds 4 million fish, management of the commercial fishery is significantly altered, including: (1) the Central District Drift Gillnet Fishery Management Plan (5 AAC 21.353), which states that there will be no mandatory restrictions to the fishery from July 16 to July 31, (2) the Northern District Salmon Management Plan (5 AAC 21.358), which says the Yentna River sockeye salmon sustainable escapement goal of 90,000 to 160,000 fish changes to an optimal escapement goal of 75,000 to 180,000 fish, and (3) the Kenai River Late-Run Sockeye Salmon Management Plan (5 AAC 21.360), which directs ADF&G to manage for an inriver goal range of 850,000 to 1,100,000 fish and requires only one continuous 36 hour closure to the Upper Subdistrict set gillnet fishery. In 2005, the official announcement that the Kenai River sockeye salmon run would likely exceed 4 million fish occurred on 26 July and the various management plan changes just outlined went into effect. The test fish data was heavily relied upon to make this formal projection.

The last day of test fishing typically occurs on 30 July each year, which means the “tail-end” of the sockeye salmon run is not assessed by the project. Two methods were developed to estimate the percentage of the run that occurred after the test fishery ceased so that postseason adjustments could be made to the *CCPUE* statistic to reflect what it would have been had the project continued through the end of the sockeye salmon run. The first method accounts for the number of fish harvested commercially after the test fishery ends (Equation 13), while the second method enumerates both escapement and commercial catch (total run) after the test fishery terminates (Equation 14). The sport and personal-use harvest of sockeye salmon that occurs after the test fishery concludes is assumed to be very minimal and therefore is not considered. Table 6 shows the differences in the annual *CCPUE* statistic after postseason adjustments were made using either the harvest or total run method. Although the changes are relatively minor, they do have an effect on the algorithms that are used to fit the current year's *CCPUE* to run-timing curves from previous years, because the a and b coefficients in the equation describing the historical run timings are changed. Beginning in 2002, the total run method was used to make postseason adjustments to all previous years' *CCPUE* statistics (Shields 2003). For the 2005 season, the final test fish *CCPUE* of 2,643 was adjusted to 3,180; in other words, approximately 20.3% of the total sockeye salmon run occurred after the test fishery ceased.

A non-linear mathematical model (Mundy 1979) was fit to the *CCPUE* proportions of the sockeye salmon run to UCI. Using the total run-adjusted *CCPUE*, this analysis suggested that 6.4% of the run had passed the transect prior to the start of test fishing on 1 July and that the run was 75.2% complete at project termination (Figure 3 and Appendix A13). Therefore, the mathematical model indicated that test fishing spanned only 68.8% of the run. The median date

of the 2005 UCI sockeye salmon run (the day on which approximately 50% of the total run has entered UCI at the test fish transect) occurred on day 28.4, or 21 July, which was 6 days late relative to the historic average median date of 15 July (Table 7). This entry timing represents the latest run that had ever been observed since the test fish project began in 1979. Moreover, 39% of the sockeye salmon escapement into the Kenai River occurred in August, which is the second largest percentage of escapement in August since escapement monitoring began in 1978. Incidentally, the midpoint date of the sockeye salmon run at the test fish transect in 1994 was 20 July (Table 7), the second latest midpoint date of run since the test fishery began.

The strength of the latter portion of the 2005 run, however, was predicted inseason by the test fish program. For example, from 21 July to 30 July, the test fish *CCPUE* was 1,360 points, which suggested that as many as 3.7 million sockeye salmon had entered the district during this time period. The cumulative run through July 21 was estimated at 4.4 million (Table 4) fish, while the total run in 2005 was estimated at 7.8 million, so about 3.4 million fish went into catch and escapement after July 20. This is only slightly below (0.1 million) what the *OTF CCPUE* had suggested. In addition, 2.7 million sockeye salmon were harvested after July 20, which is slightly more than half of the total harvest. Surprisingly though, there was no single significant harvest event within the first week after the large test fish catches, instead the later harvests were more spread out over time.

Water temperatures measured along the test fish transect ranged from 9.6° to 14.5°C and averaged 11.7°C for the year (Appendices A14 and A15). The average surface water temperature was the warmest observed at each station during the past 10 years. Strangely, these warmer than average surface water temperatures did not fit with the run-timing of this year's run. Burgner (1980) showed that the arrival dates of Bristol Bay sockeye salmon were early during years where water temperatures were warmer than average, while Blackbourn (1987) provided a model that explained the negative relationship between sea surface temperatures in the Gulf of Alaska and the timing of the run of sockeye salmon stocks from Alaska. Therefore, based on surface water temperatures at the test fish transect in 2005, which were warmer than average, it could have been expected that this year's run would have been early. In the third volume of a series of books dealing with Fishery Oceanography, Pearcy (1992) summarized many of the factors that affect the coastal migration of returning adult salmon. He reviewed the orientation mechanisms used by salmon in coastal waters and concluded that prior to entering estuaries, adult salmon probably rely on cues that are different from those used in the open ocean phases of their migration. Salinity, temperature, currents, and bathymetry were all thought to play a role in migration. Another factor to consider that could affect run timing is the age composition of the run, which relates to fish size; larger fish swim faster than smaller fish (Flynn and Hilborn 2004). It is altogether possible that some combination of these factors (other than just water temperature) may affect fish migration and run timing.

For the past 2 years, ADF&G has been conducting a companion study on board the test fish vessel, using side-scan sonar to observe fish distribution in the water column in relation to various oceanographic data, such as water temperature, salinity, tide stage, and water clarity. These data have not been published yet, but one of the objectives of the study was to determine whether or not the *OTF* inseason run forecasting model could be enhanced and improved using this additional information.

Air temperatures along the test fish transect ranged from 10° to 19°C and averaged 12.9°C, with station averages nearly identical to the previous 10-year averages (Appendices A14 and A15).

Wind velocities averaged 7.1 knots for the month, which represents the calmest conditions observed during the past 10 years. Wind direction was variable, but in general winds originated out of the south. The average water and air temperature from all stations combined was warmer than long-term values, while the average wind velocity was less than long-term measurements (Appendix 16). Salinity and secchi disk readings were nearly identical to the averages from all previous years.

Appendix A15 provides a summary of the physical data that has been collected at each of the 6 test fish stations for the past 10 years. Station 4, which is on the east side of Cook Inlet, was the shallowest station, averaging 24.0 fathoms (144 feet) in depth. It should be noted that changes in depth are a result of different stages of tide as well as minor differences in set location from day to day. Station 4 also had the clearest water, with a 1996–2005 secchi disk average depth of 8.1 m. In general, water clarity along the test fish transect decreases as you travel from east to west (secchi disk average depth decreases from 8.1 m at station 4 to 2.8 m at station 8) as a result of numerous glacial watersheds draining into the west side of Cook Inlet.

Since 2002, scale samples have been collected from all sockeye salmon that are measured to estimate mean length. The dominant age-class of sockeye salmon entering UCI at the test fish transect in 2005 was 1.3 (5-year olds), comprising nearly 80% of the run (Table 8). This estimate compared favorably with age composition samples collected from sockeye salmon harvested commercially by the drift gillnet fleet in UCI, which revealed the 1.3 age class comprised 79% of the 2005 harvest (T. Tobias, Commercial Fisheries Technician, ADF&G, Soldotna; personal communication). The scale samples from the OTF program were also analyzed with the intent of assessing whether or not Kenai River sockeye salmon, which are the dominant stock in Cook Inlet runs, might be identified using “size at age” criteria as they entered the district at the test fish transect. Statistical analyses will be conducted comparing the average size of each age class of sockeye salmon collected at various escapement monitoring sites throughout UCI to the average size of the same age classes collected at the 6 stations along the test fish transect. The results of this “mixture-model” analysis will be summarized in future test fish annual reports.

The UCI test fishery continues to provide fishery managers with very important data about the strength and timing of each year’s sockeye salmon run. And, because commercial, sport, and personal use fishery management plans are intrinsically linked to the strength of the annual sockeye salmon run, the UCI test fishery is essentially relied upon to provide fishermen and fishery managers with an early indication of annual run strength and timing.

ACKNOWLEDGEMENTS

The authors would like to acknowledge Roy Self, captain of the *FV Corrina Kay*, for his expertise in conducting safe and efficient maritime activities. After the 2005 season, Captain Self retired from skippering the test fish vessel after 27 consecutive years of service. The Alaska Department of Fish and Game would like to express our sincerest gratitude to Captain Self, as he demonstrated a commitment to the integrity of the test fish project that was very admirable and appreciated. Captain Self was always cordial and often made very helpful suggestions about ways to improve the project. He also was keenly aware that the data being collected was important to advancing the understanding and management of UCI salmon stocks. He played an integral role in the development and success of the project. We wish him all the best in his future endeavors.

In addition, we would like to thank Scott Pegau (Kachemak Bay National Estuarine Research Reserve) for his help in test fish data collection while he was on board the *FV Corrina Kay* conducting oceanographic studies.

REFERENCES CITED

- Clutter, R., and L. Whitesel. 1956. Collection and interpretation of sockeye salmon scales. Bulletin International Pacific Salmon Fisheries Commission No. 9, New Westminster, B.C.
- Blackbourn, D. J. 1987. Sea surface temperature and pre-season prediction of return timing in Fraser River sockeye salmon (*Oncorhynchus nerka*). Pages 269-306 [In]: Population biology and future management, Smith, H. D., L. Margolis, and C. C. Wood, editors. Canadian Special Publication of Fisheries and Aquatic Sciences 96.
- Burgner, R. L. 1980. Some features of ocean migrations and timing of Pacific salmon. Pages 153-164 [In]: Salmonid ecosystems of the North Pacific, McNeil, W., and D. C. Himsworth, editors. Corvallis: Oregon State University Press.
- Flynn L., and R. Hilborn. 2004. Test fishery indices for sockeye salmon (*Oncorhynchus nerka*) as affected by age composition and environmental variables. Canadian Journal of Fisheries and Aquatic Sciences 61(1):80-92.
- Hilsinger, J. R. 1988. Run strength analysis of the 1987 sockeye salmon return to Upper Cook Inlet, Alaska. Alaska Department of Fish and Game, Division of Commercial Fisheries, Regional Information Report 2A88-19, Anchorage.
- Hilsinger, J. R., and D. Waltemyer. 1987. Run strength analysis of the 1986 sockeye salmon return to Upper Cook Inlet, Alaska. Alaska Department of Fish and Game, Division of Commercial Fisheries, Upper Cook Inlet Area Data Report 87-6, Soldotna.
- Koo, T. S. Y. 1955. Biology of the red salmon, *Oncorhynchus nerka* (Walbaum), of Bristol Bay, Alaska as revealed by a study of their scales. Doctoral dissertation, University of Washington, Seattle.
- Koo, T. S. Y. 1962. Age determination in salmon. Pages 37-48 [In]: Studies of Alaska red salmon, T. S. Y. Koo, editor. University of Washington Press, Seattle.
- Mosher, K. 1969. Identification of Pacific salmon and steelhead trout by scale characteristics. United States Department of the Interior, Circular 317, Washington, D.C.
- Mundy, P. R. 1979. A quantitative measure of migratory timing illustrated by application to the management of commercial salmon fisheries. Doctoral dissertation, University of Washington, Seattle, Washington.
- Mundy, P. R., K. K. English, W. J. Gazey, and K. E. Tarbox. 1993. Evaluation of the harvest management strategies applied to sockeye salmon populations of Upper Cook Inlet, Alaska, using run reconstruction analysis. [In]: Kruse G., D. M. Eggers, R. J. Marasco, C. Pautzke, and T. J. Quinn II editors. Proceedings of the international symposium on management strategies for exploited fish populations. Alaska Sea Grant College Program, University of Alaska, Fairbanks.
- Pearcy, W. G. 1992. Ocean ecology of North Pacific salmonids. Washington Sea Grant Program. University of Washington Press, Seattle, WA.
- Shields, P. 2006. Upper Cook Inlet commercial fisheries annual management report. 2005. Alaska Department of Fish and Game, Fishery Management Report No. 06-42, Anchorage.
<http://www.sf.adfg.state.ak.us/FedAidPDFs/fmr06-42.pdf>
- Shields, P. A. 2000. An estimate of the migratory timing and abundance of sockeye salmon into Upper Cook Inlet, Alaska, in 2000. Alaska Department of Fish and Game, Commercial Fisheries Management and Development Division, Regional Information Report 2A00-30, Anchorage.
- Shields, P. A. 2001. An estimate of the migratory timing and abundance of sockeye salmon into Upper Cook Inlet, Alaska, in 2001. Alaska Department of Fish and Game, Division of Commercial Fisheries, Regional Information Report 2A01-14, Anchorage.

REFERENCES CITED (Continued)

- Shields, P. A. 2003. An estimate of the migratory timing and abundance of sockeye salmon into Upper Cook Inlet, Alaska, in 2002. Alaska Department of Fish and Game, Division of Commercial Fisheries, Regional Information Report 2A03-01, Anchorage.
- Shields, P. A., and T. M. Willette. 2004. An estimate of the migratory timing and abundance of sockeye salmon into Upper Cook Inlet, Alaska, in 2003. Alaska Department of Fish and Game, Division of Commercial Fisheries, Regional Information Report 2A04-15, Anchorage.
- Shields, P., and M. Willette. 2005. An estimate of the migratory timing and abundance of sockeye salmon into Upper Cook Inlet, Alaska, 2004. Alaska Department of Fish and Game, Fishery Data Series No. 05-64, Anchorage. <http://www.sf.adfg.state.ak.us/FedAidPDFs/fds05-64.pdf>
- Tarbox, K. E. 1990. An estimate of the migratory timing of sockeye salmon into Upper Cook Inlet, Alaska, in 1989 using a test fishery. Alaska Department of Fish and Game, Division of Commercial Fisheries, Regional Information Report 2A90-4, Anchorage.
- Tarbox, K. E. 1991. An estimate of the migratory timing of sockeye salmon into Upper Cook, Alaska, in 1990 using a test fishery. Alaska Department of Fish and Game, Division of Commercial Fisheries, Regional Information Report 2A91-6, Anchorage.
- Tarbox, K. E. 1994. An estimate of the migratory timing and abundance of sockeye salmon into Upper Cook Inlet, Alaska, in 1993. Alaska Department of Fish and Game, Commercial Fisheries Management and Development Division, Regional Information Report 2A94-13, Anchorage.
- Tarbox, K. E. 1995. An estimate of migratory timing and abundance of sockeye salmon into Upper Cook Inlet, Alaska, in 1994. Alaska Department of Fish and Game, Commercial Fisheries Management and Development Division, Regional Information Report 2A95-15, Anchorage.
- Tarbox, K. E. 1996. An estimate of migratory timing and abundance of sockeye salmon into Upper Cook Inlet, Alaska, in 1995. Alaska Department of Fish and Game, Commercial Fisheries Management and Development Division, Regional Information Report 2A96-07, Anchorage.
- Tarbox, K. E. 1997. An estimate of migratory timing and abundance of sockeye salmon into Upper Cook Inlet, Alaska, in 1996. Alaska Department of Fish and Game, Commercial Fisheries Management and Development Division, Regional Information Report 2A97-01, Anchorage.
- Tarbox, K. E. 1998a. An estimate of migratory timing and abundance of sockeye salmon into Upper Cook Inlet, Alaska, in 1997. Alaska Department of Fish and Game, Commercial Fisheries Management and Development Division, Regional Information Report 2A98-22, Anchorage.
- Tarbox, K. E. 1998b. An estimate of migratory timing and abundance of sockeye salmon into Upper Cook Inlet, Alaska, in 1998. Alaska Department of Fish and Game, Commercial Fisheries Management and Development Division, Regional Information Report 2A98-30, Anchorage.
- Tarbox, K. E. 1999. An estimate of migratory timing and abundance of sockeye salmon into Upper Cook Inlet, Alaska, in 1999. Alaska Department of Fish and Game, Commercial Fisheries Management and Development Division, Regional Information Report 2A99-13, Anchorage.
- Tarbox, K. E., and B. King 1992. An estimate of the migratory timing of sockeye salmon into Upper Cook Inlet, Alaska, in 1991 using a test fishery. Regional Information Report 2A92-07, Anchorage.
- Tarbox, K. E., and D. Waltemyer. 1989. An estimate of the 1988 total sockeye salmon return to Upper Cook Inlet, Alaska using a test fishery. Alaska Department of Fish and Game, Division of Commercial Fisheries, Regional Information Report 2A89-4, Anchorage.
- Tobias, T. M., D. L. Waltemyer, and K. E. Tarbox. 1994. Scale aging manual for Upper Cook Inlet sockeye salmon. Alaska Department of Fish and Game, Division of Commercial Fisheries Management and Development, Regional Information Report 2A94-36, Anchorage.

REFERENCES CITED (Continued)

- Tobias, T. M., and M. Willette. 2003. An estimate of total return of sockeye salmon to Upper Cook Inlet, Alaska, 1976-2002. Alaska Department of Fish and Game, Division of Commercial Fisheries, Regional Information Report 2A03-11, Anchorage.
- Waltemyer, D. L. 1983a. Migratory timing and abundance estimation of the 1982 sockeye salmon return to Upper Cook Inlet based on a test fishing program. Alaska Department of Fish and Game, Division of Commercial Fisheries, Upper Cook Inlet Data Report 83-1, Soldotna.
- Waltemyer, D. L. 1983b. Describing the migrations of salmon and estimating abundance of sockeye salmon returning in 1983 to Upper Cook Inlet based on a test fishery. Alaska Department of Fish and Game, Division of Commercial Fisheries, Upper Cook Inlet Data Report 84-1, Soldotna.
- Waltemyer, D. L. 1986a. Use of a test fishery to describe and estimate the sockeye salmon total return to Upper Cook Inlet in 1984. Alaska Department of Fish and Game, Division of Commercial Fisheries, Upper Cook Inlet Data Report 86-1, Soldotna.
- Waltemyer, D. L. 1986b. Run strength analysis of the 1985 sockeye salmon return to Upper Cook Inlet, Alaska based on a test fishery. Alaska Department of Fish and Game, Division of Commercial Fisheries, Upper Cook Inlet Data Report 86-5, Soldotna.

TABLES AND FIGURES

Table 1.–Summary of sockeye salmon fishing effort, daily and cumulative catch, and daily and cumulative CPUE, Upper Cook Inlet offshore test fish project, 2005.

| Date | Number of Stations | Mean Fishing Time (min) | CATCH | | | | Mean Length (mm) |
|-------------------|--------------------|-------------------------|-------|-------|-------|-------|------------------|
| | | | CATCH | | CPUE | | |
| | | | Daily | Cum | Daily | Cum | |
| 7/01 | 6 | 231.5 | 47 | 47 | 35 | 35 | 556 |
| 7/02 | 6 | 230.0 | 43 | 90 | 33 | 68 | 544 |
| 7/03 | 6 | 240.0 | 142 | 232 | 98 | 165 | 559 |
| 7/04 | 6 | 227.5 | 46 | 278 | 36 | 201 | 560 |
| 7/05 | 6 | 239.7 | 68 | 346 | 47 | 249 | 568 |
| 7/06 | 6 | 241.0 | 94 | 440 | 68 | 317 | 563 |
| 7/07 | 6 | 246.5 | 171 | 611 | 118 | 434 | 565 |
| 7/08 | 6 | 256.0 | 196 | 807 | 127 | 561 | 566 |
| 7/09 | 6 | 252.5 | 199 | 1,006 | 124 | 685 | 572 |
| 7/10 ^a | 6 | 236.2 | 66 | 1,072 | 46 | 731 | 569 |
| 7/11 | 6 | 235.0 | 50 | 1,122 | 36 | 767 | 568 |
| 7/12 | 6 | 233.5 | 81 | 1,203 | 60 | 827 | 565 |
| 7/13 | 6 | 235.5 | 100 | 1,303 | 76 | 903 | 569 |
| 7/14 | 6 | 225.5 | 38 | 1,341 | 30 | 932 | 568 |
| 7/15 | 6 | 267.0 | 217 | 1,558 | 136 | 1,068 | 572 |
| 7/16 | 6 | 252.5 | 35 | 1,593 | 25 | 1,093 | 578 |
| 7/17 | 6 | 227.0 | 39 | 1,632 | 30 | 1,122 | 570 |
| 7/18 | 6 | 234.0 | 56 | 1,688 | 37 | 1,160 | 577 |
| 7/19 | 6 | 220.0 | 37 | 1,725 | 29 | 1,188 | 570 |
| 7/20 | 6 | 245.5 | 149 | 1,874 | 95 | 1,283 | 572 |
| 7/21 | 6 | 267.5 | 294 | 2,168 | 168 | 1,451 | 576 |
| 7/22 | 6 | 242.0 | 136 | 2,304 | 94 | 1,545 | 565 |
| 7/23 | 6 | 257.0 | 212 | 2,516 | 133 | 1,678 | 569 |
| 7/24 | 6 | 243.0 | 128 | 2,644 | 91 | 1,769 | 565 |
| 7/25 | 6 | 256.0 | 238 | 2,882 | 153 | 1,922 | 567 |
| 7/26 | 6 | 258.0 | 227 | 3,109 | 149 | 2,071 | 570 |
| 7/27 | 6 | 222.0 | 25 | 3,134 | 19 | 2,090 | 567 |
| 7/28 | 6 | 243.5 | 196 | 3,330 | 148 | 2,238 | 572 |
| 7/29 | 6 | 260.0 | 311 | 3,641 | 196 | 2,435 | 575 |
| 7/30 | 6 | 287.0 | 467 | 4,108 | 208 | 2,643 | 571 |

^a Two of the 6 stations were not fished due to weather; the data for these stations were interpolated.

Table 2.—Estimated sockeye salmon catch by date and station, Upper Cook Inlet offshore test fish project, 2005.

| Date | Station Number | | | | | | Total |
|-------------------|----------------|--------------|--------------|--------------|--------------|--------------|--------------|
| | 4 | 5 | 6 | 6.5 | 7 | 8 | |
| 7/01 | 0 | 5 | 36 | 3 | 2 | 1 | 47 |
| 7/02 | 1 | 15 | 11 | 12 | 4 | 0 | 43 |
| 7/03 | 50 | 7 | 1 | 73 | 10 | 1 | 142 |
| 7/04 | 3 | 4 | 13 | 22 | 3 | 1 | 46 |
| 7/05 | 0 | 6 | 13 | 43 | 6 | 0 | 68 |
| 7/06 | 5 | 22 | 18 | 4 | 44 | 1 | 94 |
| 7/07 | 15 | 6 | 29 | 72 | 48 | 1 | 171 |
| 7/08 | 7 | 93 | 21 | 39 | 34 | 2 | 196 |
| 7/09 | 0 | 1 | 34 | 69 | 94 | 1 | 199 |
| 7/10 ^a | 0 | 14 | 39 | 12 | 1 | 0 | 66 |
| 7/11 | 1 | 27 | 2 | 0 | 3 | 17 | 50 |
| 7/12 | 0 | 25 | 3 | 32 | 8 | 13 | 81 |
| 7/13 | 13 | 2 | 16 | 16 | 28 | 25 | 100 |
| 7/14 | 0 | 7 | 2 | 8 | 3 | 18 | 38 |
| 7/15 | 12 | 20 | 17 | 103 | 43 | 22 | 217 |
| 7/16 | 3 | 1 | 9 | 2 | 12 | 8 | 35 |
| 7/17 | 8 | 4 | 1 | 2 | 24 | 0 | 39 |
| 7/18 | 6 | 38 | 9 | 1 | 2 | 0 | 56 |
| 7/19 | 4 | 6 | 1 | 5 | 3 | 18 | 37 |
| 7/20 | 3 | 11 | 97 | 35 | 0 | 3 | 149 |
| 7/21 | 1 | 16 | 16 | 45 | 48 | 168 | 294 |
| 7/22 | 2 | 26 | 7 | 1 | 59 | 41 | 136 |
| 7/23 | 1 | 0 | 18 | 63 | 91 | 39 | 212 |
| 7/24 | 6 | 14 | 18 | 13 | 20 | 57 | 128 |
| 7/25 | 1 | 3 | 91 | 26 | 72 | 45 | 238 |
| 7/26 | 0 | 31 | 76 | 72 | 22 | 26 | 227 |
| 7/27 | 0 | 0 | 0 | 18 | 2 | 5 | 25 |
| 7/28 | 20 | 69 | 39 | 31 | 31 | 6 | 196 |
| 7/29 | 19 | 91 | 47 | 131 | 23 | 0 | 311 |
| 7/30 | 12 | 10 | 338 | 87 | 19 | 1 | 467 |
| Total | 193 | 574 | 1,022 | 1,040 | 759 | 520 | 4,108 |
| Percent | 4.7% | 14.0% | 24.9% | 25.3% | 18.5% | 12.7% | 100% |

^a Two of the 6 stations were not fished due to weather; the data for these stations were interpolated.

Table 3.—Estimated sockeye salmon CPUE by date and station, Upper Cook Inlet offshore test fish project, 2005.

| Date | Station Number | | | | | | Total |
|-------------------|----------------|--------------|--------------|--------------|--------------|--------------|--------------|
| | 4 | 5 | 6 | 6.5 | 7 | 8 | |
| 7/01 | 0 | 4 | 26 | 2 | 2 | 1 | 35 |
| 7/02 | 1 | 12 | 8 | 9 | 3 | 0 | 33 |
| 7/03 | 36 | 6 | 1 | 47 | 8 | 1 | 98 |
| 7/04 | 2 | 3 | 10 | 17 | 2 | 1 | 36 |
| 7/05 | 0 | 5 | 9 | 29 | 5 | 0 | 47 |
| 7/06 | 4 | 16 | 14 | 3 | 31 | 1 | 68 |
| 7/07 | 12 | 5 | 19 | 50 | 32 | 1 | 118 |
| 7/08 | 5 | 55 | 15 | 26 | 23 | 2 | 127 |
| 7/09 | 0 | 1 | 24 | 41 | 57 | 1 | 124 |
| 7/10 ^a | 0 | 10 | 26 | 9 | 1 | 0 | 46 |
| 7/11 | 1 | 19 | 2 | 0 | 2 | 13 | 36 |
| 7/12 | 0 | 19 | 2 | 22 | 6 | 10 | 60 |
| 7/13 | 10 | 2 | 12 | 12 | 20 | 19 | 76 |
| 7/14 | 0 | 5 | 2 | 6 | 2 | 14 | 30 |
| 7/15 | 9 | 15 | 13 | 56 | 28 | 15 | 136 |
| 7/16 | 2 | 1 | 6 | 2 | 9 | 6 | 25 |
| 7/17 | 6 | 3 | 1 | 2 | 18 | 0 | 30 |
| 7/18 | 5 | 23 | 7 | 1 | 2 | 0 | 37 |
| 7/19 | 3 | 5 | 1 | 4 | 2 | 13 | 29 |
| 7/20 | 2 | 9 | 58 | 24 | 0 | 2 | 95 |
| 7/21 | 1 | 12 | 12 | 31 | 32 | 81 | 168 |
| 7/22 | 2 | 19 | 6 | 1 | 39 | 28 | 94 |
| 7/23 | 1 | 0 | 13 | 39 | 54 | 26 | 133 |
| 7/24 | 5 | 11 | 13 | 10 | 15 | 38 | 91 |
| 7/25 | 1 | 2 | 54 | 20 | 44 | 33 | 153 |
| 7/26 | 0 | 21 | 47 | 46 | 16 | 19 | 149 |
| 7/27 | 0 | 0 | 0 | 14 | 2 | 4 | 19 |
| 7/28 | 15 | 56 | 27 | 23 | 23 | 4 | 148 |
| 7/29 | 14 | 56 | 31 | 77 | 18 | 0 | 196 |
| 7/30 | 9 | 8 | 125 | 51 | 14 | 1 | 208 |
| Total | 146 | 399 | 582 | 674 | 510 | 331 | 2,643 |
| Percent | 5.5% | 15.1% | 22.0% | 25.5% | 19.3% | 12.5% | 100% |

^a Two of the 6 stations were not fished due to weather; the data for these stations were interpolated.

Table 4.–Total run estimates for sockeye salmon to Upper Cook Inlet, Alaska, made during the 2005 season.

| Based on data through 7/21/2005 | | | | | | |
|---|------------------------|----------------------|--------------|------------|--------------|------------------------|
| Escapement | | | | | | 1,158,782 |
| Cumulative Catch | | | | | | 2,909,242 |
| Residual in District | | | | | | 298,693 |
| Total Run Through 7/21/2005 = | | | | | | 4,366,718 |
| 2005 Cumulative OTF CPUE through 7/21/2005 = | | | | | | 1,451 |
| Passage Rate (Total Run/Cumulative CPUE) based on 7/21/2005 harvest = | | | | | | 3,009 |
| Run Estimates Based on Model Results (Fit of Current Year to Past Years) | | | | | | |
| Year | Mean Sum of Squares | Estimated Total CPUE | | | Timing | Estimated Total Run |
| | | Current | Previous Day | Difference | | |
| 1994 | 0.001599 | 3,048 | 3,099 | -51 | Late 4 days | 9,171,546 |
| 1997 | 0.001746 | 2,451 | 2,471 | -19 | Late 1 day | 7,375,241 |
| 1998 | 0.001786 | 2,414 | 2,433 | -19 | Late 3 days | 7,263,442 |
| 1985 | 0.001924 | 1,987 | 1,986 | 1 | On Time | 5,976,972 |
| 1988 | 0.001947 | 1,921 | 1,918 | 3 | Early 2 days | 5,780,151 |
| 1982 | 0.001956 | 2,116 | 2,124 | -8 | Late 2 days | 6,366,764 |
| 2001 | 0.001982 | 1,651 | 1,642 | 9 | Early 2 days | 4,967,382 |
| 2002 | 0.002086 | 1,665 | 1,658 | 7 | Early 2 days | 5,008,900 |
| 1986 | 0.002146 | 2,076 | 2,087 | -11 | Late 1 day | 6,246,090 |
| 1993 | 0.002195 | 1,956 | 1,962 | -6 | Early 1 day | 5,885,571 |
| 1984 | 0.002629 | 1,574 | 1,554 | 20 | Early 4 days | 4,735,751 |
| 1983 | 0.003049 | 2,119 | 2,143 | -24 | On Time | 6,375,820 |
| 1987 | 0.003287 | 2,879 | 2,962 | -83 | Late 2 days | 8,661,381 |
| 1996 | 0.003321 | 1,809 | 1,817 | -8 | Early 2 days | 5,442,467 |
| 2003 | 0.003455 | 1,913 | 1,928 | -15 | Early 1 day | 5,756,232 |
| 2004 | 0.003917 | 2,481 | 2,539 | -58 | Late 2 days | 7,463,724 |
| 1991 | 0.003975 | 2,430 | 2,485 | -55 | Late 2 days | 7,309,504 |
| 1989 | 0.004116 | 1,973 | 1,950 | 24 | On Time | 5,936,206 |
| 1995 | 0.004202 | 2,045 | 2,073 | -28 | On Time | 6,152,162 |
| 2000 | 0.004219 | 1,653 | 1,654 | -2 | Early 2 days | 4,971,804 |
| 1979 | 0.004614 | 1,398 | 1,369 | 29 | Early 5 days | 4,206,150 |
| 1999 | 0.005580 | 2,626 | 2,716 | -90 | Late 3 days | 7,899,728 |
| 1990 | 0.006694 | 3,028 | 3,181 | -153 | Late 3 days | 9,108,877 |
| 1992 | 0.006838 | 2,550 | 2,644 | -94 | Late 2 days | 7,671,797 |
| 1981 | 0.019323 | 1,264 | 1,225 | 39 | Early 9 days | 3,803,120 |
| 1980 | 0.021614 | 1,298 | 1,259 | 40 | Early 9 days | 3,905,352 |

-continued-

Table 4.—Page 2 of 2.

| Based on data through 7/25/2005 | | | | | | |
|---|------------------------|----------------------|--------------|------------|--------------|------------------------|
| Escapement | | | | | | 1,401,796 |
| Cumulative Catch | | | | | | 3,650,920 |
| Residual in District | | | | | | 269,511 |
| TOTAL RUN THROUGH 7/25/2005 = | | | | | | 5,322,227 |
| 2004 Cumulative OTF CPUE through 7/25/2005 = | | | | | | 1,922 |
| Passage Rate (Total Run/Cumulative CPUE) through 7/25/2005 = | | | | | | 2,769 |
| Run Estimates Based on Model Results (Fit of Current Year to Past Years) | | | | | | |
| Year | Mean Sum of Squares | Estimated Total CPUE | | | Timing | Estimated Total Run |
| | | Current | Previous Day | Difference | | |
| 1994 | 0.001531 | 2,947 | 2,954 | -7 | Late 4 days | 8,161,541 |
| 1997 | 0.001541 | 2,459 | 2,442 | 17 | Late 1 day | 6,809,306 |
| 1998 | 0.001583 | 2,424 | 2,407 | 18 | Late 3 days | 6,713,954 |
| 1982 | 0.001942 | 2,165 | 2,138 | 26 | Late 2 days | 5,995,229 |
| 1986 | 0.002102 | 2,120 | 2,094 | 26 | Late 1 day | 5,870,023 |
| 1985 | 0.002280 | 2,064 | 2,031 | 33 | On Time | 5,716,319 |
| 1993 | 0.002378 | 2,017 | 1,987 | 30 | Early 1 day | 5,585,823 |
| 1988 | 0.002516 | 2,007 | 1,973 | 35 | Early 2 days | 5,559,264 |
| 1983 | 0.002736 | 2,127 | 2,108 | 19 | On Time | 5,890,960 |
| 2003 | 0.003344 | 1,955 | 1,928 | 27 | Early 2 days | 5,413,757 |
| 1987 | 0.003480 | 2,701 | 2,722 | -21 | Late 2 days | 7,480,258 |
| 1996 | 0.003605 | 1,873 | 1,841 | 31 | Early 2 days | 5,186,414 |
| 2004 | 0.003712 | 2,386 | 2,388 | -2 | Late 2 days | 6,607,137 |
| 1991 | 0.003714 | 2,346 | 2,346 | 0 | Late 2 days | 6,496,996 |
| 1995 | 0.003751 | 2,048 | 2,029 | 19 | On Time | 5,672,174 |
| 2002 | 0.003918 | 1,772 | 1,732 | 40 | Early 1 days | 4,907,559 |
| 2001 | 0.004058 | 1,763 | 1,722 | 41 | Early 2 days | 4,883,603 |
| 1989 | 0.005235 | 2,119 | 2,072 | 47 | On Time | 5,867,835 |
| 2000 | 0.005303 | 1,741 | 1,704 | 37 | Early 2 days | 4,821,430 |
| 1999 | 0.005743 | 2,450 | 2,467 | -17 | Late 3 days | 6,783,938 |
| 1984 | 0.006357 | 1,717 | 1,670 | 47 | Early 4 days | 4,755,849 |
| 1992 | 0.006961 | 2,373 | 2,389 | -16 | Late 2 days | 6,571,799 |
| 1990 | 0.008017 | 2,683 | 2,733 | -49 | Late 3 days | 7,431,655 |
| 1979 | 0.012549 | 1,569 | 1,516 | 53 | Early 5 days | 4,346,498 |
| 1981 | 0.032592 | 1,466 | 1,406 | 59 | Early 9 days | 4,059,529 |
| 1980 | 0.033135 | 1,501 | 1,442 | 60 | Early 9 days | 4,157,235 |

Table 5.—Estimated total Kenai River sockeye run (millions) in 2005 estimated from total offshore test fish CPUE and age composition stock allocation estimates through July 21, and July 27.

| Data through 21-July | | | | | | | | | | | | |
|-----------------------------|------------|----------------------------|------------------|---------------|---------------------|----------------------|------------------------|--------------------------|--------------------------|--------------------|----------------------------|------------------|
| Year | MSS | Est. Total OTF CPUE | | | Passage Rate | Estimated | Estimated | Estimated | Estimated | Prop. Kenai | Estimated | Estimated |
| | | Current | Prev. Day | Timing | | UCI Total run | UCI Run to Date | UCI Run Remaining | Kenai Run to Date | | Kenai Run Remaining | Kenai Run |
| 1994 | 0.00160 | 3,048 | 3,099 | Late 4 days | 3,009 | 9.2 | 4.1 | 5.1 | 2.6 | 65% | 3.3 | 5.9 |
| 1997 | 0.00175 | 2,451 | 2,471 | Late 1 day | 3,009 | 7.4 | 4.1 | 3.3 | 2.6 | 65% | 2.1 | 4.7 |
| 1998 | 0.00179 | 2,414 | 2,433 | Late 3 days | 3,009 | 7.3 | 4.1 | 3.2 | 2.6 | 65% | 2.1 | 4.6 |
| 1985 | 0.00192 | 1,987 | 1,986 | On Time | 3,009 | 6.0 | 4.1 | 1.9 | 2.6 | 65% | 1.2 | 3.8 |
| 1988 | 0.00195 | 1,921 | 1,918 | Early 2 days | 3,009 | 5.8 | 4.1 | 1.7 | 2.6 | 65% | 1.1 | 3.7 |

| Data through 25-July | | | | | | | | | | | | |
|-----------------------------|------------|----------------------------|------------------|---------------|---------------------|----------------------|------------------------|--------------------------|--------------------------|--------------------|----------------------------|------------------|
| Year | MSS | Est. Total OTF CPUE | | | Passage Rate | Estimated | Estimated | Estimated | Estimated | Prop. Kenai | Estimated | Estimated |
| | | Current | Prev. Day | Timing | | UCI Total run | UCI Run to Date | UCI Run Remaining | Kenai Run to Date | | Kenai Run Remaining | Kenai Run |
| 1994 | 0.00153 | 2,947 | 2,954 | Late 4 days | 2,769 | 8.2 | 5.1 | 3.1 | 3.3 | 69% | 2.1 | 5.5 |
| 1997 | 0.00154 | 2,459 | 2,442 | Late 1 day | 2,769 | 6.8 | 5.1 | 1.8 | 3.3 | 69% | 1.2 | 4.6 |
| 1998 | 0.00158 | 2,424 | 2,407 | Late 3 days | 2,769 | 6.7 | 5.1 | 1.7 | 3.3 | 69% | 1.1 | 4.5 |
| 1982 | 0.00194 | 2,165 | 2,138 | Late 2 days | 2,769 | 6.0 | 5.1 | 0.9 | 3.3 | 69% | 0.6 | 4.0 |
| 1986 | 0.00210 | 2,120 | 2,094 | Late 1 day | 2,769 | 5.9 | 5.1 | 0.8 | 3.3 | 69% | 0.6 | 3.9 |

Note: MSS is the mean sum of squares.

Table 6.—A comparison of methods used to make post-season adjustments to the offshore test fish final CPUE.

| Year | Final OTF CPUE | Post-Season OTF CPUE Adjustment | | Harvest Adjusted | | Total Run Adjusted | |
|------|----------------|---------------------------------|--------------------|------------------|----------|--------------------|----------|
| | | Harvest-adjusted | Total Run-adjusted | <i>a</i> | <i>b</i> | <i>a</i> | <i>b</i> |
| 1979 | 602 | 651 | 664 | -3.2451 | 0.1876 | -3.3380 | 0.2004 |
| 1980 | 740 | 770 | 777 | -2.2537 | 0.1640 | -2.2403 | 0.1612 |
| 1981 | 364 | 383 | 387 | -2.5459 | 0.1856 | -2.5243 | 0.1819 |
| 1982 | 651 | 775 | 786 | -3.6839 | 0.1522 | -3.7156 | 0.1633 |
| 1983 | 2,464 | 2,472 | 2,474 | -4.2719 | 0.1883 | -4.2732 | 0.1884 |
| 1984 | 1,331 | 1,334 | 1,341 | -3.4257 | 0.1855 | -3.4018 | 0.1834 |
| 1985 | 1,422 | 1,575 | 1,563 | -3.4581 | 0.1523 | -3.5633 | 0.1626 |
| 1986 | 1,653 | 1,731 | 1,714 | -3.7671 | 0.1633 | -3.8642 | 0.1719 |
| 1987 | 1,404 | 1,422 | 1,428 | -4.3442 | 0.1689 | -4.6385 | 0.1785 |
| 1988 | 1,131 | 1,145 | 1,169 | -3.3682 | 0.1639 | -3.5655 | 0.1662 |
| 1989 | 619 | 682 | 692 | -2.7114 | 0.1258 | -2.7031 | 0.1238 |
| 1990 | 1,358 | 1,404 | 1,426 | -5.7913 | 0.2259 | -5.7085 | 0.2211 |
| 1991 | 1,574 | 1,759 | 1,740 | -4.5806 | 0.1885 | -4.6331 | 0.1919 |
| 1992 | 2,021 | 2,186 | 2,195 | -5.4366 | 0.2235 | -5.4043 | 0.2217 |
| 1993 | 1,815 | 1,882 | 1,913 | -4.0776 | 0.1906 | -3.9018 | 0.1797 |
| 1994 | 1,012 | 1,145 | 1,199 | -4.0770 | 0.1553 | -3.9757 | 0.1453 |
| 1995 | 1,712 | 1,828 | 1,850 | -4.7036 | 0.2131 | -4.6219 | 0.2078 |
| 1996 | 1,723 | 1,765 | 1,796 | -4.6328 | 0.2266 | -4.4605 | 0.2144 |
| 1997 | 1,656 | 1,705 | 1,826 | -3.8265 | 0.1621 | -3.7000 | 0.1496 |
| 1998 | 1,158 | 1,355 | 1,313 | -3.6700 | 0.1473 | -3.7142 | 0.1515 |
| 1999 | 2,226 | 2,475 | 2,419 | -5.3100 | 0.2175 | -5.1500 | 0.2081 |
| 2000 | 1,520 | 1,532 | 1,565 | -5.1094 | 0.2614 | -4.9141 | 0.2480 |
| 2001 | 1,586 | 1,594 | 1,630 | -3.9323 | 0.2002 | -3.9823 | 0.2041 |
| 2002 | 1,736 | 1,749 | 1,825 | -4.3694 | 0.2292 | -4.0642 | 0.2068 |
| 2003 | 1,787 | 1,824 | 1,848 | -4.5091 | 0.2117 | -4.4402 | 0.2068 |
| 2004 | 2,028 | 2,220 | 2,345 | -4.6374 | 0.1903 | -4.6374 | 0.1903 |
| 2005 | 2,643 | 3,015 | 3,180 | -3.7499 | 0.1360 | -3.7170 | 0.1305 |

Table 7.—Mean date of the sockeye salmon run across the Anchor Point transect, Upper Cook Inlet offshore test fish project, 1979–2005.

| Year | Mean Date ^a | |
|-------------------|------------------------|----------|
| | Coded | Calendar |
| 1979 | 16.7 | 10-Jul |
| 1980 | 13.9 | 7-Jul |
| 1981 | 13.9 | 7-Jul |
| 1982 | 22.8 | 16-Jul |
| 1983 | 22.7 | 16-Jul |
| 1984 | 18.5 | 12-Jul |
| 1985 | 21.9 | 15-Jul |
| 1986 | 22.5 | 16-Jul |
| 1987 | 26.0 | 19-Jul |
| 1988 | 21.4 | 14-Jul |
| 1989 | 21.8 | 15-Jul |
| 1990 | 25.8 | 19-Jul |
| 1991 | 24.1 | 17-Jul |
| 1992 | 24.4 | 17-Jul |
| 1993 | 21.7 | 15-Jul |
| 1994 | 27.4 | 20-Jul |
| 1995 | 22.2 | 15-Jul |
| 1996 | 20.8 | 14-Jul |
| 1997 | 24.7 | 18-Jul |
| 1998 | 24.5 | 18-Jul |
| 1999 | 24.7 | 18-Jul |
| 2000 | 19.8 | 13-Jul |
| 2001 | 19.5 | 13-Jul |
| 2002 | 19.7 | 13-Jul |
| 2003 | 21.5 | 14-Jul |
| 2004 | 24.4 | 17-Jul |
| 1979–2004 Average | 21.8 | 15-Jul |
| 2005 | 28.4 | 21-Jul |

^a Day (1) = June 24.

Table 8.—Sockeye salmon age-composition, mean length (mm), and number of samples by station, Upper Cook Inlet offshore test fish project, 2005.

| Station No. | Sample Size | Percent Age Composition | | | | | | | | |
|----------------|-------------|-------------------------|-----|------|-----|-----|------|-----|-----|-----|
| | | 0.3 | 1.2 | 1.3 | 2.2 | 1.4 | 2.3 | 2.4 | 3.2 | 3.3 |
| 4 | 126 | | 2.6 | 85.5 | 1.0 | 1.0 | 8.8 | | 1.0 | |
| 5 | 319 | 0.9 | 1.9 | 82.6 | 1.1 | 0.7 | 12.9 | | | |
| 6 | 361 | 0.6 | 1.4 | 80.0 | 1.4 | 0.3 | 16.1 | 0.3 | | |
| 6.5 | 439 | 0.5 | 3.3 | 78.9 | 1.5 | 0.4 | 14.8 | 0.2 | 0.2 | 0.2 |
| 7 | 392 | | 1.8 | 78.0 | 1.6 | 1.3 | 17.3 | | | |
| 8 | 244 | 0.8 | 1.2 | 77.5 | 2.5 | 0.4 | 17.7 | | | |
| Weighted Avg = | | 0.7 | 2.3 | 79.9 | 1.7 | 0.9 | 15.6 | 0.2 | 0.6 | 0.2 |

| Station No. | Sample Size | Mean Length (mm) | | | | | | | | |
|----------------|-------------|------------------|-----|-----|-----|-----|-----|-----|-----|-----|
| | | 0.3 | 1.2 | 1.3 | 2.2 | 1.4 | 2.3 | 2.4 | 3.2 | 3.3 |
| 4 | 126 | | 489 | 570 | 521 | 594 | 563 | | 511 | |
| 5 | 319 | 570 | 494 | 571 | 515 | 602 | 572 | | | |
| 6 | 361 | 564 | 496 | 570 | 503 | 581 | 570 | 621 | | |
| 6.5 | 439 | 568 | 506 | 572 | 509 | 608 | 568 | 592 | 542 | 561 |
| 7 | 392 | | 522 | 575 | 531 | 554 | 561 | | | |
| 8 | 244 | 585 | 534 | 571 | 528 | 629 | 564 | | | |
| Weighted Avg = | | 572 | 507 | 572 | 518 | 583 | 567 | 607 | 527 | 561 |

| Station No. | Sample Size | Number of Samples | | | | | | | | |
|-----------------|-------------|-------------------|-----|-------|-----|-----|-----|-----|-----|-----|
| | | 0.3 | 1.2 | 1.3 | 2.2 | 1.4 | 2.3 | 2.4 | 3.2 | 3.3 |
| 4 | 126 | | 3 | 109 | 1 | 1 | 11 | | 1 | |
| 5 | 319 | 3 | 6 | 264 | 3 | 2 | 41 | | | |
| 6 | 361 | 2 | 5 | 289 | 5 | 1 | 58 | 1 | | |
| 6.5 | 439 | 2 | 14 | 346 | 7 | 2 | 65 | 1 | 1 | 1 |
| 7 | 392 | | 7 | 306 | 6 | 5 | 68 | | | |
| 8 | 244 | 2 | 3 | 189 | 6 | 1 | 43 | | | |
| Station Total = | | 9 | 38 | 1,503 | 28 | 12 | 286 | 2 | 2 | 1 |




Figure 1.—Location of fishing districts and offshore test fish transect in Cook Inlet, Alaska 2005.




Figure 2.—Linear regression of the relationship between offshore test fish unadjusted cumulative CPUE and Upper Cook Inlet sockeye salmon total annual run.




Figure 3.—Cumulative proportions estimated for the sockeye salmon run to Upper Cook Inlet, Alaska, 2005.




Figure 4.—OTF error in forecasting the total run of sockeye salmon to Upper Cook Inlet using the 20-July best-fit estimate.

APPENDIX A.

Appendix A1.–Summary of pink salmon fishing effort, daily and cumulative catch, and daily and cumulative CPUE, Upper Cook Inlet offshore test fish project, 2005.

| Date | Number of Stations | Mean Fishing Time (min) | CATCH | | CPUE | |
|-------------------|---------------------------|--------------------------------|--------------|------------|--------------|------------|
| | | | Daily | Cum | Daily | Cum |
| 7/01 | 6 | 231.5 | 1 | 1 | 1 | 1 |
| 7/02 | 6 | 230.0 | 3 | 4 | 2 | 3 |
| 7/03 | 6 | 240.0 | 4 | 8 | 3 | 6 |
| 7/04 | 6 | 227.5 | 4 | 12 | 3 | 9 |
| 7/05 | 6 | 239.7 | 3 | 15 | 2 | 11 |
| 7/06 | 6 | 241.0 | 3 | 18 | 2 | 13 |
| 7/07 | 6 | 246.5 | 4 | 22 | 3 | 16 |
| 7/08 | 6 | 256.0 | 14 | 36 | 9 | 25 |
| 7/09 | 6 | 252.5 | 8 | 44 | 5 | 31 |
| 7/10 ^a | 6 | 236.2 | 14 | 58 | 11 | 41 |
| 7/11 | 6 | 235.0 | 11 | 69 | 8 | 50 |
| 7/12 | 6 | 233.5 | 11 | 80 | 9 | 58 |
| 7/13 | 6 | 235.5 | 7 | 87 | 5 | 63 |
| 7/14 | 6 | 225.5 | 2 | 89 | 2 | 65 |
| 7/15 | 6 | 267.0 | 11 | 100 | 27 | 92 |
| 7/16 | 6 | 252.5 | 5 | 105 | 4 | 95 |
| 7/17 | 6 | 227.0 | 4 | 109 | 3 | 99 |
| 7/18 | 6 | 234.0 | 0 | 109 | 0 | 99 |
| 7/19 | 6 | 220.0 | 2 | 111 | 2 | 100 |
| 7/20 | 6 | 245.5 | 6 | 117 | 4 | 104 |
| 7/21 | 6 | 267.5 | 10 | 127 | 5 | 110 |
| 7/22 | 6 | 242.0 | 6 | 133 | 4 | 114 |
| 7/23 | 6 | 257.0 | 5 | 138 | 3 | 117 |
| 7/24 | 6 | 243.0 | 10 | 148 | 7 | 124 |
| 7/25 | 6 | 256.0 | 10 | 158 | 7 | 131 |
| 7/26 | 6 | 258.0 | 4 | 162 | 3 | 134 |
| 7/27 | 6 | 222.0 | 3 | 165 | 2 | 136 |
| 7/28 | 6 | 243.5 | 11 | 176 | 8 | 144 |
| 7/29 | 6 | 260.0 | 5 | 181 | 3 | 148 |
| 7/30 | 6 | 287.0 | 5 | 186 | 2 | 150 |

^a Two of the 6 stations were not fished due to weather; the data for these stations were interpolated.

Appendix A2.—Estimated pink salmon catch by date and station, Upper Cook Inlet offshore test fish project, 2005.

| Date | Station Number | | | | | | Total |
|-------------------|----------------|------|------|------|------|------|-------|
| | 4 | 5 | 6 | 6.5 | 7 | 8 | |
| 7/01 | 0 | 0 | 1 | 0 | 0 | 0 | 1 |
| 7/02 | 0 | 2 | 0 | 0 | 0 | 1 | 3 |
| 7/03 | 1 | 1 | 0 | 2 | 0 | 0 | 4 |
| 7/04 | 0 | 1 | 1 | 0 | 2 | 0 | 4 |
| 7/05 | 0 | 0 | 1 | 2 | 0 | 0 | 3 |
| 7/06 | 1 | 1 | 1 | 0 | 0 | 0 | 3 |
| 7/07 | 1 | 0 | 2 | 1 | 0 | 0 | 4 |
| 7/08 | 2 | 5 | 1 | 3 | 3 | 0 | 14 |
| 7/09 | 0 | 1 | 0 | 0 | 6 | 1 | 8 |
| 7/10 ^a | 2 | 1 | 6 | 4 | 0 | 1 | 14 |
| 7/11 | 5 | 2 | 0 | 1 | 2 | 1 | 11 |
| 7/12 | 0 | 1 | 5 | 1 | 1 | 3 | 11 |
| 7/13 | 1 | 1 | 3 | 2 | 0 | 0 | 7 |
| 7/14 | 0 | 0 | 0 | 0 | 1 | 1 | 2 |
| 7/15 | 0 | 2 | 2 | 4 | 2 | 1 | 11 |
| 7/16 | 0 | 0 | 1 | 1 | 2 | 1 | 5 |
| 7/17 | 0 | 0 | 0 | 1 | 2 | 1 | 4 |
| 7/18 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 7/19 | 0 | 0 | 1 | 0 | 1 | 0 | 2 |
| 7/20 | 2 | 0 | 4 | 0 | 0 | 0 | 6 |
| 7/21 | 1 | 0 | 0 | 1 | 0 | 8 | 10 |
| 7/22 | 0 | 0 | 0 | 0 | 1 | 5 | 6 |
| 7/23 | 0 | 0 | 1 | 2 | 2 | 0 | 5 |
| 7/24 | 0 | 2 | 6 | 0 | 0 | 2 | 10 |
| 7/25 | 1 | 1 | 2 | 2 | 1 | 3 | 10 |
| 7/26 | 0 | 0 | 0 | 3 | 1 | 0 | 4 |
| 7/27 | 0 | 0 | 1 | 1 | 1 | 0 | 3 |
| 7/28 | 3 | 0 | 1 | 0 | 5 | 2 | 11 |
| 7/29 | 1 | 0 | 3 | 1 | 0 | 0 | 5 |
| 7/30 | 0 | 0 | 4 | 1 | 0 | 0 | 5 |
| Total | 21 | 21 | 47 | 33 | 33 | 31 | 186 |
| Percent | 11.3 | 11.3 | 25.3 | 17.7 | 17.7 | 16.7 | 100 |

^a Two of the 6 stations were not fished due to weather; the data for these stations were interpolated.

Appendix A3.—Estimated pink salmon CPUE by date and station, Upper Cook Inlet offshore test fish project, 2005.

| Date | Station Number | | | | | | Total |
|-------------------|----------------|-------------|-------------|-------------|-------------|-------------|--------------|
| | 4 | 5 | 6 | 6.5 | 7 | 8 | |
| 7/01 | 0.0 | 0.0 | 0.7 | 0.0 | 0.0 | 0.0 | 0.7 |
| 7/02 | 0.0 | 1.5 | 0.0 | 0.0 | 0.0 | 0.8 | 2.3 |
| 7/03 | 0.7 | 0.8 | 0.0 | 1.3 | 0.0 | 0.0 | 2.8 |
| 7/04 | 0.0 | 0.8 | 0.8 | 0.0 | 1.6 | 0.0 | 3.2 |
| 7/05 | 0.0 | 0.0 | 0.7 | 1.4 | 0.0 | 0.0 | 2.1 |
| 7/06 | 0.7 | 0.7 | 0.7 | 0.0 | 0.0 | 0.0 | 2.1 |
| 7/07 | 0.8 | 0.0 | 1.3 | 0.7 | 0.0 | 0.0 | 2.8 |
| 7/08 | 1.6 | 2.9 | 0.7 | 2.0 | 2.0 | 0.0 | 9.2 |
| 7/09 | 0.0 | 0.8 | 0.0 | 0.0 | 3.7 | 0.8 | 5.3 |
| 7/10 ^a | 1.9 | 1.1 | 4.0 | 3.0 | 0.0 | 0.8 | 10.8 |
| 7/11 | 3.8 | 1.4 | 0.0 | 0.8 | 1.5 | 0.7 | 8.3 |
| 7/12 | 0.0 | 0.7 | 3.9 | 0.7 | 0.8 | 2.3 | 8.5 |
| 7/13 | 0.8 | 0.8 | 2.2 | 1.5 | 0.0 | 0.0 | 5.4 |
| 7/14 | 0.0 | 0.0 | 0.0 | 0.0 | 0.8 | 0.8 | 1.6 |
| 7/15 | 0.0 | 1.5 | 1.5 | 21.8 | 1.3 | 0.6 | 26.7 |
| 7/16 | 0.0 | 0.0 | 0.6 | 0.9 | 1.5 | 0.7 | 3.7 |
| 7/17 | 0.0 | 0.0 | 0.0 | 0.8 | 1.5 | 0.8 | 3.1 |
| 7/18 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| 7/19 | 0.0 | 0.0 | 1.1 | 0.0 | 0.7 | 0.0 | 1.8 |
| 7/20 | 1.6 | 0.0 | 2.4 | 0.0 | 0.0 | 0.0 | 4.0 |
| 7/21 | 0.8 | 0.0 | 0.0 | 0.7 | 0.0 | 3.8 | 5.4 |
| 7/22 | 0.0 | 0.0 | 0.0 | 0.0 | 0.7 | 3.4 | 4.1 |
| 7/23 | 0.0 | 0.0 | 0.7 | 1.2 | 1.2 | 0.0 | 3.2 |
| 7/24 | 0.0 | 1.5 | 4.4 | 0.0 | 0.0 | 1.3 | 7.3 |
| 7/25 | 0.8 | 0.8 | 1.2 | 1.5 | 0.6 | 2.2 | 7.1 |
| 7/26 | 0.0 | 0.0 | 0.0 | 1.9 | 0.7 | 0.0 | 2.6 |
| 7/27 | 0.0 | 0.0 | 0.9 | 0.7 | 0.8 | 0.0 | 2.4 |
| 7/28 | 2.2 | 0.0 | 0.7 | 0.0 | 3.6 | 1.4 | 8.0 |
| 7/29 | 0.7 | 0.0 | 2.0 | 0.6 | 0.0 | 0.0 | 3.3 |
| 7/30 | 0.0 | 0.0 | 1.5 | 0.6 | 0.0 | 0.0 | 2.1 |
| Total | 16.5 | 15.3 | 32.1 | 42.1 | 23.1 | 20.5 | 149.7 |
| Percent | 11.0 | 10.2 | 21.4 | 28.2 | 15.5 | 13.7 | 100 |

^a Two of the 6 stations were not fished due to weather; the data for these stations were interpolated.

Appendix A4.–Summary of chum salmon fishing effort, daily and cumulative catch, and daily and cumulative CPUE, Upper Cook Inlet offshore test fish project, 2005.

| Date | Number of Stations | Mean Fishing Time (min) | CATCH | | CPUE | |
|-------------------|--------------------|-------------------------|-------|-----|-------|-----|
| | | | Daily | Cum | Daily | Cum |
| 7/01 | 6 | 231.5 | 2 | 2 | 2 | 2 |
| 7/02 | 6 | 230.0 | 2 | 4 | 2 | 3 |
| 7/03 | 6 | 240.0 | 6 | 10 | 4 | 7 |
| 7/04 | 6 | 227.5 | 2 | 12 | 2 | 8 |
| 7/05 | 6 | 239.7 | 5 | 17 | 3 | 12 |
| 7/06 | 6 | 241.0 | 16 | 33 | 12 | 23 |
| 7/07 | 6 | 246.5 | 17 | 50 | 12 | 35 |
| 7/08 | 6 | 256.0 | 37 | 87 | 24 | 59 |
| 7/09 | 6 | 252.5 | 21 | 108 | 13 | 73 |
| 7/10 ^a | 6 | 236.2 | 19 | 127 | 13 | 86 |
| 7/11 | 6 | 235.0 | 12 | 139 | 8 | 94 |
| 7/12 | 6 | 233.5 | 11 | 150 | 8 | 102 |
| 7/13 | 6 | 235.5 | 6 | 156 | 5 | 107 |
| 7/14 | 6 | 225.5 | 3 | 159 | 3 | 110 |
| 7/15 | 6 | 267.0 | 55 | 214 | 33 | 142 |
| 7/16 | 6 | 252.5 | 5 | 219 | 3 | 145 |
| 7/17 | 6 | 227.0 | 8 | 227 | 6 | 152 |
| 7/18 | 6 | 234.0 | 7 | 234 | 4 | 156 |
| 7/19 | 6 | 220.0 | 6 | 240 | 4 | 160 |
| 7/20 | 6 | 245.5 | 15 | 255 | 10 | 171 |
| 7/21 | 6 | 267.5 | 35 | 290 | 22 | 193 |
| 7/22 | 6 | 242.0 | 6 | 296 | 4 | 197 |
| 7/23 | 6 | 257.0 | 22 | 318 | 14 | 211 |
| 7/24 | 6 | 243.0 | 19 | 337 | 14 | 224 |
| 7/25 | 6 | 256.0 | 21 | 358 | 14 | 238 |
| 7/26 | 6 | 258.0 | 30 | 388 | 20 | 258 |
| 7/27 | 6 | 222.0 | 4 | 392 | 3 | 261 |
| 7/28 | 6 | 243.5 | 22 | 414 | 17 | 278 |
| 7/29 | 6 | 260.0 | 17 | 431 | 11 | 289 |
| 7/30 | 6 | 287.0 | 17 | 448 | 11 | 300 |

^a Two of the 6 stations were not fished due to weather; the data for these stations were interpolated.

Appendix A5.—Estimated chum salmon catch by date and station, Upper Cook Inlet offshore test fish project, 2005.

| Date | Station Number | | | | | | Total |
|-------------------|----------------|------------|------------|------------|------------|------------|-------------|
| | 4 | 5 | 6 | 6.5 | 7 | 8 | |
| 7/01 | 0 | 0 | 0 | 2 | 0 | 0 | 2 |
| 7/02 | 0 | 1 | 0 | 1 | 0 | 0 | 2 |
| 7/03 | 0 | 0 | 0 | 6 | 0 | 0 | 6 |
| 7/04 | 0 | 0 | 1 | 0 | 1 | 0 | 2 |
| 7/05 | 0 | 0 | 0 | 5 | 0 | 0 | 5 |
| 7/06 | 0 | 1 | 2 | 1 | 11 | 1 | 16 |
| 7/07 | 0 | 0 | 3 | 4 | 8 | 2 | 17 |
| 7/08 | 0 | 11 | 2 | 14 | 8 | 2 | 37 |
| 7/09 | 0 | 0 | 5 | 5 | 11 | 0 | 21 |
| 7/10 ^a | 0 | 6 | 8 | 3 | 1 | 1 | 19 |
| 7/11 | 0 | 12 | 0 | 0 | 0 | 0 | 12 |
| 7/12 | 0 | 1 | 6 | 4 | 0 | 0 | 11 |
| 7/13 | 0 | 2 | 1 | 3 | 0 | 0 | 6 |
| 7/14 | 0 | 0 | 0 | 0 | 3 | 0 | 3 |
| 7/15 | 0 | 2 | 1 | 33 | 17 | 2 | 55 |
| 7/16 | 1 | 0 | 3 | 0 | 1 | 0 | 5 |
| 7/17 | 0 | 0 | 0 | 1 | 7 | 0 | 8 |
| 7/18 | 0 | 6 | 1 | 0 | 0 | 0 | 7 |
| 7/19 | 0 | 0 | 0 | 0 | 1 | 5 | 6 |
| 7/20 | 0 | 5 | 8 | 1 | 1 | 0 | 15 |
| 7/21 | 0 | 5 | 3 | 6 | 10 | 11 | 35 |
| 7/22 | 0 | 2 | 1 | 0 | 0 | 3 | 6 |
| 7/23 | 0 | 0 | 4 | 6 | 9 | 3 | 22 |
| 7/24 | 0 | 1 | 3 | 5 | 3 | 7 | 19 |
| 7/25 | 0 | 0 | 2 | 0 | 9 | 10 | 21 |
| 7/26 ^a | 0 | 3 | 17 | 4 | 2 | 4 | 30 |
| 7/27 | 0 | 1 | 2 | 1 | 0 | 0 | 4 |
| 7/28 | 1 | 12 | 2 | 2 | 4 | 1 | 22 |
| 7/29 | 2 | 6 | 4 | 5 | 0 | 0 | 17 |
| 7/30 | 0 | 0 | 1 | 8 | 7 | 1 | 17 |
| Total | 4 | 77 | 80 | 120 | 114 | 53 | 448 |
| Percent | 1% | 17% | 18% | 27% | 25% | 12% | 100% |

^a Two of the 6 stations were not fished due to weather; the data for these stations were interpolated.

Appendix A6.—Estimated chum salmon CPUE by date and station, Upper Cook Inlet offshore test fish project, 2005.

| Date | Station Number | | | | | | Total |
|-------------------|----------------|-------------|-------------|-------------|-------------|-------------|--------------|
| | 4 | 5 | 6 | 6.5 | 7 | 8 | |
| 7/01 | 0.0 | 0.0 | 0.0 | 1.5 | 0.0 | 0.0 | 1.5 |
| 7/02 | 0.0 | 0.7 | 0.0 | 0.8 | 0.0 | 0.0 | 1.5 |
| 7/03 | 0.0 | 0.0 | 0.0 | 3.8 | 0.0 | 0.0 | 3.8 |
| 7/04 | 0.0 | 0.0 | 0.8 | 0.0 | 0.8 | 0.0 | 1.6 |
| 7/05 | 0.0 | 0.0 | 0.0 | 3.4 | 0.0 | 0.0 | 3.4 |
| 7/06 | 0.0 | 0.7 | 1.5 | 0.8 | 7.7 | 0.8 | 11.5 |
| 7/07 | 0.0 | 0.0 | 2.0 | 2.8 | 5.2 | 1.6 | 11.6 |
| 7/08 | 0.0 | 6.5 | 1.5 | 9.4 | 5.4 | 1.6 | 24.4 |
| 7/09 | 0.0 | 0.0 | 3.5 | 3.0 | 6.7 | 0.0 | 13.2 |
| 7/10 ^a | 0.0 | 4.1 | 5.3 | 2.3 | 0.8 | 0.8 | 13.3 |
| 7/11 | 0.0 | 8.3 | 0.0 | 0.0 | 0.0 | 0.0 | 8.3 |
| 7/12 | 0.0 | 0.7 | 4.7 | 2.8 | 0.0 | 0.0 | 8.3 |
| 7/13 | 0.0 | 1.6 | 0.7 | 2.3 | 0.0 | 0.0 | 4.6 |
| 7/14 | 0.0 | 0.0 | 0.0 | 0.0 | 2.5 | 0.0 | 2.5 |
| 7/15 | 0.0 | 1.5 | 0.7 | 18.0 | 11.1 | 1.3 | 32.6 |
| 7/16 | 0.8 | 0.0 | 1.9 | 0.0 | 0.7 | 0.0 | 3.4 |
| 7/17 | 0.0 | 0.0 | 0.0 | 0.8 | 5.2 | 0.0 | 6.0 |
| 7/18 | 0.0 | 3.7 | 0.8 | 0.0 | 0.0 | 0.0 | 4.4 |
| 7/19 | 0.0 | 0.0 | 0.0 | 0.0 | 0.7 | 3.7 | 4.4 |
| 7/20 | 0.0 | 3.9 | 4.7 | 0.7 | 0.8 | 0.0 | 10.1 |
| 7/21 | 0.0 | 3.7 | 2.2 | 4.1 | 6.7 | 5.3 | 22.0 |
| 7/22 | 0.0 | 1.4 | 0.8 | 0.0 | 0.0 | 2.0 | 4.2 |
| 7/23 | 0.0 | 0.0 | 3.0 | 3.7 | 5.3 | 2.0 | 14.0 |
| 7/24 | 0.0 | 0.7 | 2.2 | 3.8 | 2.2 | 4.6 | 13.6 |
| 7/25 | 0.0 | 0.0 | 1.2 | 0.0 | 5.5 | 7.2 | 13.9 |
| 7/26 ^a | 0.0 | 2.1 | 10.5 | 2.5 | 1.4 | 2.9 | 19.5 |
| 7/27 | 0.0 | 0.8 | 1.8 | 0.7 | 0.0 | 0.0 | 3.3 |
| 7/28 | 0.7 | 9.7 | 1.4 | 1.5 | 2.9 | 0.7 | 17.0 |
| 7/29 | 1.5 | 3.7 | 2.6 | 2.9 | 0.0 | 0.0 | 10.8 |
| 7/30 | 0.0 | 0.0 | 0.4 | 4.7 | 5.3 | 0.8 | 11.2 |
| Total | 3.0 | 53.9 | 54.2 | 76.4 | 77.0 | 35.3 | 300.0 |
| Percent | 1 | 18 | 18 | 26 | 26 | 12 | 100 |

^a Two of the 6 stations were not fished due to weather; the data for these stations were interpolated.

Appendix A7.—Summary of coho salmon fishing effort, daily and cumulative catch, and daily and cumulative CPUE, Upper Cook Inlet offshore test fish project, 2005.

| Date | Number of Stations | Mean Fishing Time (min) | CATCH | | CPUE | |
|-------------------|--------------------|-------------------------|-------|-----|-------|-----|
| | | | Daily | Cum | Daily | Cum |
| 7/01 | 6 | 231.5 | 3 | 3 | 2 | 2 |
| 7/02 | 6 | 230.0 | 0 | 3 | 0 | 2 |
| 7/03 | 6 | 240.0 | 4 | 7 | 3 | 5 |
| 7/04 | 6 | 227.5 | 0 | 7 | 0 | 5 |
| 7/05 | 6 | 239.7 | 1 | 8 | 1 | 6 |
| 7/06 | 6 | 241.0 | 0 | 8 | 0 | 6 |
| 7/07 | 6 | 246.5 | 2 | 10 | 1 | 7 |
| 7/08 | 6 | 256.0 | 14 | 24 | 10 | 17 |
| 7/09 | 6 | 252.5 | 1 | 25 | 1 | 18 |
| 7/10 ^a | 6 | 236.2 | 7 | 32 | 5 | 23 |
| 7/11 | 6 | 235.0 | 0 | 32 | 0 | 23 |
| 7/12 | 6 | 233.5 | 3 | 35 | 2 | 25 |
| 7/13 | 6 | 235.5 | 12 | 47 | 9 | 34 |
| 7/14 | 6 | 225.5 | 1 | 48 | 1 | 35 |
| 7/15 | 6 | 267.0 | 21 | 69 | 14 | 49 |
| 7/16 | 6 | 252.5 | 3 | 72 | 2 | 51 |
| 7/17 | 6 | 227.0 | 5 | 77 | 4 | 55 |
| 7/18 | 6 | 234.0 | 8 | 85 | 5 | 60 |
| 7/19 | 6 | 220.0 | 2 | 87 | 2 | 61 |
| 7/20 | 6 | 245.5 | 14 | 101 | 10 | 71 |
| 7/21 | 6 | 267.5 | 55 | 156 | 29 | 100 |
| 7/22 | 6 | 242.0 | 7 | 163 | 5 | 105 |
| 7/23 | 6 | 257.0 | 57 | 220 | 36 | 141 |
| 7/24 | 6 | 243.0 | 37 | 257 | 27 | 168 |
| 7/25 | 6 | 256.0 | 67 | 324 | 42 | 210 |
| 7/26 | 6 | 258.0 | 25 | 349 | 17 | 227 |
| 7/27 | 6 | 222.0 | 7 | 356 | 6 | 233 |
| 7/28 | 6 | 243.5 | 30 | 386 | 23 | 255 |
| 7/29 | 6 | 260.0 | 81 | 467 | 51 | 307 |
| 7/30 | 6 | 287.0 | 79 | 546 | 37 | 344 |

^a Two of the 6 stations were not fished due to weather; the data for these stations were interpolated.

Appendix A8.—Estimated coho salmon catch by date and station, Upper Cook Inlet offshore test fish project, 2005.

| Date | Station Number | | | | | | Total |
|-------------------|----------------|------------|------------|------------|------------|------------|-------------|
| | 4 | 5 | 6 | 6.5 | 7 | 8 | |
| 7/01 | 1 | 1 | 1 | 0 | 0 | 0 | 3 |
| 7/02 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 7/03 | 3 | 0 | 0 | 1 | 0 | 0 | 4 |
| 7/04 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 7/05 | 0 | 0 | 0 | 1 | 0 | 0 | 1 |
| 7/06 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 7/07 | 0 | 0 | 1 | 1 | 0 | 0 | 2 |
| 7/08 | 0 | 2 | 7 | 0 | 4 | 1 | 14 |
| 7/09 | 0 | 0 | 0 | 0 | 1 | 0 | 1 |
| 7/10 ^a | 0 | 0 | 4 | 1 | 2 | 0 | 7 |
| 7/11 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 7/12 | 0 | 0 | 2 | 1 | 0 | 0 | 3 |
| 7/13 | 1 | 2 | 6 | 0 | 2 | 1 | 12 |
| 7/14 | 0 | 0 | 0 | 1 | 0 | 0 | 1 |
| 7/15 | 1 | 7 | 2 | 5 | 3 | 3 | 21 |
| 7/16 | 0 | 0 | 2 | 0 | 1 | 0 | 3 |
| 7/17 | 0 | 1 | 0 | 0 | 4 | 0 | 5 |
| 7/18 | 1 | 5 | 1 | 0 | 0 | 1 | 8 |
| 7/19 | 0 | 0 | 0 | 0 | 0 | 2 | 2 |
| 7/20 | 0 | 1 | 5 | 1 | 0 | 7 | 14 |
| 7/21 | 0 | 0 | 5 | 2 | 3 | 45 | 55 |
| 7/22 | 0 | 0 | 2 | 0 | 2 | 3 | 7 |
| 7/23 | 0 | 0 | 5 | 14 | 21 | 17 | 57 |
| 7/24 | 0 | 2 | 10 | 6 | 16 | 3 | 37 |
| 7/25 | 0 | 0 | 11 | 2 | 49 | 5 | 67 |
| 7/26 ^a | 0 | 0 | 6 | 4 | 7 | 8 | 25 |
| 7/27 | 0 | 0 | 2 | 2 | 2 | 1 | 7 |
| 7/28 | 0 | 8 | 5 | 13 | 3 | 1 | 30 |
| 7/29 | 2 | 23 | 29 | 24 | 0 | 3 | 81 |
| 7/30 | 1 | 8 | 53 | 13 | 3 | 1 | 79 |
| Total | 10 | 60 | 159 | 92 | 123 | 102 | 546 |
| Percent | 2% | 11% | 29% | 17% | 23% | 19% | 100% |

^a Two of the 6 stations were not fished due to weather; the data for these stations were interpolated.

Appendix A9.—Estimated coho salmon CPUE by date and station, Upper Cook Inlet offshore test fish project, 2005.

| Date | Station Number | | | | | | Total |
|-------------------|----------------|------|------|------|------|------|-------|
| | 4 | 5 | 6 | 6.5 | 7 | 8 | |
| 7/01 | 0.8 | 0.8 | 0.7 | 0.0 | 0.0 | 0.0 | 2.3 |
| 7/02 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| 7/03 | 2.1 | 0.0 | 0.0 | 0.6 | 0.0 | 0.0 | 2.7 |
| 7/04 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| 7/05 | 0.0 | 0.0 | 0.0 | 0.7 | 0.0 | 0.0 | 0.7 |
| 7/06 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| 7/07 | 0.0 | 0.0 | 0.7 | 0.7 | 0.0 | 0.0 | 1.4 |
| 7/08 | 0.0 | 1.2 | 5.1 | 0.0 | 2.7 | 0.8 | 9.8 |
| 7/09 | 0.0 | 0.0 | 0.0 | 0.0 | 0.6 | 0.0 | 0.6 |
| 7/10 ^a | 0.0 | 0.0 | 2.6 | 0.8 | 1.6 | 0.0 | 5.0 |
| 7/11 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| 7/12 | 0.0 | 0.0 | 1.6 | 0.7 | 0.0 | 0.0 | 2.3 |
| 7/13 | 0.8 | 1.6 | 4.5 | 0.0 | 1.4 | 0.8 | 9.1 |
| 7/14 | 0.0 | 0.0 | 0.0 | 0.8 | 0.0 | 0.0 | 0.8 |
| 7/15 | 0.8 | 5.2 | 1.5 | 2.7 | 2.0 | 1.9 | 14.1 |
| 7/16 | 0.0 | 0.0 | 1.3 | 0.0 | 0.8 | 0.0 | 2.0 |
| 7/17 | 0.0 | 0.8 | 0.0 | 0.0 | 3.0 | 0.0 | 3.8 |
| 7/18 | 0.8 | 3.1 | 0.8 | 0.0 | 0.0 | 0.8 | 5.4 |
| 7/19 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 1.5 | 1.5 |
| 7/20 | 0.0 | 0.8 | 3.0 | 0.7 | 0.0 | 5.5 | 9.9 |
| 7/21 | 0.0 | 0.0 | 3.7 | 1.4 | 2.0 | 21.6 | 28.7 |
| 7/22 | 0.0 | 0.0 | 1.6 | 0.0 | 1.4 | 2.0 | 5.0 |
| 7/23 | 0.0 | 0.0 | 3.7 | 8.8 | 12.4 | 11.2 | 36.0 |
| 7/24 | 0.0 | 1.5 | 7.4 | 4.6 | 11.9 | 2.0 | 27.3 |
| 7/25 | 0.0 | 0.0 | 6.5 | 1.5 | 30.0 | 3.6 | 41.6 |
| 7/26 ^a | 0.0 | 0.0 | 3.7 | 2.6 | 5.1 | 5.8 | 17.1 |
| 7/27 | 0.0 | 0.0 | 1.8 | 1.5 | 1.6 | 0.8 | 5.7 |
| 7/28 | 0.0 | 6.5 | 3.5 | 9.6 | 2.2 | 0.7 | 22.5 |
| 7/29 | 1.5 | 14.2 | 19.1 | 14.1 | 0.0 | 2.5 | 51.4 |
| 7/30 | 0.8 | 6.2 | 19.6 | 7.6 | 2.3 | 0.8 | 37.3 |
| Total | 7.5 | 41.9 | 92.2 | 59.3 | 80.7 | 62.4 | 343.9 |
| Percent | 2 | 12 | 27 | 17 | 24 | 18 | 100 |

^a Two of the 6 stations were not fished due to weather; the data for these stations were interpolated.

Appendix A10.—Summary of Chinook salmon fishing effort, daily and cumulative catch, and daily and cumulative CPUE, Upper Cook Inlet offshore test fish project, 2005.

| Date | Number of Stations | Mean Fishing Time (min) | CATCH | | CPUE | |
|-------------------|--------------------|-------------------------|-------|-----|-------|-----|
| | | | Daily | Cum | Daily | Cum |
| 7/01 | 6 | 231.5 | 1 | 1 | 0.8 | 0.8 |
| 7/02 | 6 | 230.0 | 0 | 1 | 0.0 | 0.8 |
| 7/03 | 6 | 240.0 | 1 | 2 | 0.6 | 1.4 |
| 7/04 | 6 | 227.5 | 0 | 2 | 0.0 | 1.4 |
| 7/05 | 6 | 239.7 | 0 | 2 | 0.0 | 1.4 |
| 7/06 | 6 | 241.0 | 0 | 2 | 0.0 | 1.4 |
| 7/07 | 6 | 246.5 | 0 | 2 | 0.0 | 1.4 |
| 7/08 | 6 | 256.0 | 3 | 5 | 2.0 | 3.4 |
| 7/09 | 6 | 252.5 | 0 | 5 | 0.0 | 3.4 |
| 7/10 ^a | 6 | 236.2 | 0 | 5 | 0.0 | 3.4 |
| 7/11 | 6 | 235.0 | 1 | 6 | 0.8 | 4.2 |
| 7/12 | 6 | 233.5 | 0 | 6 | 0.0 | 4.2 |
| 7/13 | 6 | 235.5 | 0 | 6 | 0.0 | 4.2 |
| 7/14 | 6 | 225.5 | 0 | 6 | 0.0 | 4.2 |
| 7/15 | 6 | 267.0 | 1 | 7 | 0.7 | 4.8 |
| 7/16 | 6 | 252.5 | 0 | 7 | 0.0 | 4.8 |
| 7/17 | 6 | 227.0 | 0 | 7 | 0.0 | 4.8 |
| 7/18 | 6 | 234.0 | 0 | 7 | 0.0 | 4.8 |
| 7/19 | 6 | 220.0 | 0 | 7 | 0.0 | 4.8 |
| 7/20 | 6 | 245.5 | 0 | 7 | 0.0 | 4.8 |
| 7/21 | 6 | 267.5 | 0 | 7 | 0.0 | 4.8 |
| 7/22 | 6 | 242.0 | 0 | 7 | 0.0 | 4.8 |
| 7/23 | 6 | 257.0 | 0 | 7 | 0.0 | 4.8 |
| 7/24 | 6 | 243.0 | 0 | 7 | 0.0 | 4.8 |
| 7/25 | 6 | 256.0 | 0 | 7 | 0.0 | 4.8 |
| 7/26 ^a | 6 | 258.0 | 0 | 7 | 0.0 | 4.8 |
| 7/27 | 6 | 222.0 | 1 | 8 | 0.8 | 5.6 |
| 7/28 | 6 | 243.5 | 0 | 8 | 0.0 | 5.6 |
| 7/29 | 6 | 260.0 | 0 | 8 | 0.0 | 5.6 |
| 7/30 | 6 | 287.0 | 0 | 8 | 0.0 | 5.6 |

^a Two of the 6 stations were not fished due to weather; the data for these stations were interpolated.

Appendix A11.—Estimated Chinook salmon catch by date and station, Upper Cook Inlet offshore test fish project, 2005.

| Date | Station Number | | | | | | Total |
|-------------------|----------------|-----|-----|-----|-----|----|-------|
| | 4 | 5 | 6 | 6.5 | 7 | 8 | |
| 7/01 | 0 | 0 | 0 | 1 | 0 | 0 | 1 |
| 7/02 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 7/03 | 0 | 0 | 0 | 1 | 0 | 0 | 1 |
| 7/04 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 7/05 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 7/06 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 7/07 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 7/08 | 0 | 1 | 1 | 1 | 0 | 0 | 3 |
| 7/09 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 7/10 ^a | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 7/11 | 0 | 0 | 1 | 0 | 0 | 0 | 1 |
| 7/12 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 7/13 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 7/14 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 7/15 | 0 | 0 | 0 | 0 | 1 | 0 | 1 |
| 7/16 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 7/17 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 7/18 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 7/19 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 7/20 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 7/21 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 7/22 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 7/23 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 7/24 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 7/25 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 7/26 ^a | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 7/27 | 0 | 0 | 0 | 0 | 1 | 0 | 1 |
| 7/28 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 7/29 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 7/30 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Total | 0 | 1 | 2 | 3 | 2 | 0 | 8 |
| Percent | 0% | 13% | 25% | 38% | 25% | 0% | 1% |

^a Two of the 6 stations were not fished due to weather; the data for these stations were interpolated.

Appendix A12.—Estimated Chinook salmon CPUE by date and station, Upper Cook Inlet offshore test fish project, 2005.

| Date | Station Number | | | | | | Total |
|-------------------|----------------|-----|-----|-----|-----|-----|-------|
| | 4 | 5 | 6 | 6.5 | 7 | 8 | |
| 7/01 | 0.0 | 0.0 | 0.0 | 0.8 | 0.0 | 0.0 | 0.8 |
| 7/02 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| 7/03 | 0.0 | 0.0 | 0.0 | 0.6 | 0.0 | 0.0 | 0.6 |
| 7/04 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| 7/05 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| 7/06 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| 7/07 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| 7/08 | 0.0 | 0.6 | 0.7 | 0.7 | 0.0 | 0.0 | 2.0 |
| 7/09 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| 7/10 ^a | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| 7/11 | 0.0 | 0.0 | 0.8 | 0.0 | 0.0 | 0.0 | 0.8 |
| 7/12 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| 7/13 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| 7/14 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| 7/15 | 0.0 | 0.0 | 0.0 | 0.0 | 0.7 | 0.0 | 0.7 |
| 7/16 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| 7/17 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| 7/18 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| 7/19 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| 7/20 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| 7/21 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| 7/22 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| 7/23 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| 7/24 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| 7/25 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| 7/26 ^a | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| 7/27 | 0.0 | 0.0 | 0.0 | 0.0 | 0.8 | 0.0 | 0.8 |
| 7/28 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| 7/29 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| 7/30 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Total | 0.0 | 0.6 | 1.5 | 2.1 | 1.5 | 0.0 | 5.6 |
| Percent | 0% | 11% | 26% | 37% | 26% | 0% | 100% |

^a Two of the 6 stations were not fished due to weather; the data for these stations were interpolated.

Appendix A13.–Entry pattern of sockeye salmon into Upper Cook Inlet, Alaska, 2005 estimated from daily CPUE measured at the latitude of Anchor Point.

| Day | Date | Input y | Estimated y | Residual | Change in Input Y | Change in estimated Y |
|------------|-------------|--------------------|------------------------|-----------------|------------------------------|----------------------------------|
| 8 | 1-Jul | 0.0109 | 0.0642 | -0.0537 | | |
| 9 | 2-Jul | 0.0213 | 0.0729 | -0.0516 | 0.0123 | 0.0084 |
| 10 | 3-Jul | 0.0520 | 0.0823 | -0.0302 | 0.0060 | 0.0100 |
| 11 | 4-Jul | 0.0633 | 0.0927 | -0.0293 | 0.0040 | 0.0119 |
| 12 | 5-Jul | 0.0782 | 0.1042 | -0.0260 | 0.0129 | 0.0141 |
| 13 | 6-Jul | 0.0996 | 0.1171 | -0.0175 | 0.0187 | 0.0166 |
| 14 | 7-Jul | 0.1366 | 0.1312 | 0.0053 | 0.0150 | 0.0195 |
| 15 | 8-Jul | 0.1764 | 0.1468 | 0.0296 | 0.0107 | 0.0227 |
| 16 | 9-Jul | 0.2154 | 0.1639 | 0.0515 | 0.0393 | 0.0261 |
| 17 | 10-Jul | 0.2298 | 0.1826 | 0.0472 | 0.0482 | 0.0298 |
| 18 | 11-Jul | 0.2413 | 0.2029 | 0.0383 | 0.0586 | 0.0335 |
| 19 | 12-Jul | 0.2600 | 0.2248 | 0.0352 | 0.0394 | 0.0373 |
| 20 | 13-Jul | 0.2838 | 0.2484 | 0.0354 | 0.0940 | 0.0408 |
| 21 | 14-Jul | 0.2931 | 0.2735 | 0.0196 | 0.0295 | 0.0440 |
| 22 | 15-Jul | 0.3358 | 0.3002 | 0.0356 | 0.0045 | 0.0466 |
| 23 | 16-Jul | 0.3436 | 0.3283 | 0.0153 | 0.0800 | 0.0485 |
| 24 | 17-Jul | 0.3530 | 0.3577 | -0.0047 | 0.0152 | 0.0494 |
| 25 | 18-Jul | 0.3646 | 0.3882 | -0.0236 | 0.0753 | 0.0494 |
| 26 | 19-Jul | 0.3737 | 0.4196 | -0.0459 | 0.0107 | 0.0485 |
| 27 | 20-Jul | 0.4035 | 0.4517 | -0.0482 | 0.0399 | 0.0467 |
| 28 | 21-Jul | 0.4564 | 0.4842 | -0.0277 | 0.0104 | 0.0441 |
| 29 | 22-Jul | 0.4859 | 0.5168 | -0.0309 | 0.1063 | 0.0409 |
| 30 | 23-Jul | 0.5277 | 0.5492 | -0.0216 | 0.0232 | 0.0374 |
| 31 | 24-Jul | 0.5563 | 0.5813 | -0.0250 | 0.0264 | 0.0337 |
| 32 | 25-Jul | 0.6043 | 0.6127 | -0.0083 | 0.0227 | 0.0299 |
| 33 | 26-Jul | 0.6512 | 0.6432 | 0.0081 | 0.0228 | 0.0262 |
| 34 | 27-Jul | 0.6572 | 0.6725 | -0.0153 | 0.0229 | 0.0228 |
| 35 | 28-Jul | 0.7038 | 0.7006 | 0.0033 | 0.0084 | 0.0196 |
| 36 | 29-Jul | 0.7656 | 0.7272 | 0.0384 | 0.0094 | 0.0167 |
| 37 | 30-Jul | 0.8311 | 0.7523 | 0.0788 | 0.0047 | 0.0142 |

Appendix A14.—Chemical and physical observations made in Upper Cook Inlet, Alaska, during the 2005 offshore test fish project.

| Date | Sta | Air Temp °C | Water Temp (c) | Wind Vel. (knots) | Wind Dir | Tide Stage | Salinity (ppt) | Water Depth (f) | Secchi (m) |
|-------------|------------|-----------------------|--------------------------|-----------------------------|-----------------|-------------------|--------------------------|---------------------------|----------------------|
| 1-Jul | 4 | 10 | 9.9 | 5 | northwest | flood | 30.9 | 28 | 5.0 |
| | 5 | 11 | 9.6 | 4 | northwest | flood | 31.1 | 40 | 7.0 |
| | 6 | 12 | 11.9 | 6 | west | ebb | 28.0 | 53 | 3.0 |
| | 6.5 | 13 | 11.9 | 4 | southeast | ebb | 27.8 | 45 | 4.0 |
| | 7 | 13 | 12.1 | 5 | southwest | ebb | 27.7 | 50 | 4.0 |
| | 8 | 12 | 11.7 | 5 | southwest | ebb | 28.6 | 31 | 3.0 |
| 2-Jul | 8 | 10 | 11.2 | 9 | south | ebb | 28.9 | 30 | 3.0 |
| | 7 | 10 | 11.2 | 12 | southwest | ebb | 28.4 | 42 | 3.0 |
| | 6.5 | 11 | 11.6 | 12 | southwest | ebb | 27.8 | 44 | 3.0 |
| | 6 | 11 | 11.4 | 8 | south | low | 28.7 | 47 | 4.0 |
| | 5 | 11 | 10.4 | 5 | south | flood | 30.8 | 37 | 8.0 |
| | 4 | 11 | 10.0 | 5 | southwest | flood | 31.3 | 25 | 8.5 |
| 3-Jul | 4 | 12 | 10.2 | 7 | southwest | flood | 31.1 | 26 | 6.0 |
| | 5 | 11 | 10.1 | 6 | southwest | flood | 31.3 | 34 | 7.5 |
| | 6 | 12 | 11.0 | 8 | southwest | flood | 29.8 | 47 | 4.0 |
| | 6.5 | 12 | 12.5 | 8 | southwest | ebb | 27.5 | 42 | 4.0 |
| | 7 | 13 | 12.2 | 9 | south | ebb | 27.9 | 45 | 4.0 |
| | 8 | 12 | 11.8 | 15 | south | ebb | 28.5 | 29 | 5.0 |
| 4-Jul | 8 | 10 | 11.5 | 5 | south | ebb | 29.5 | 30 | 5.0 |
| | 7 | 11 | 11.3 | 5 | south | ebb | 28.8 | 40 | 3.0 |
| | 6.5 | 11 | 11.8 | 5 | south | ebb | 27.4 | 35 | 5.0 |
| | 6 | 14 | 12.2 | 5 | south | low | 26.7 | 41 | 3.0 |
| | 5 | 13 | 11.2 | 5 | south | flood | 29.7 | 36 | 6.0 |
| | 4 | 11 | 11.6 | 5 | south | flood | 31.1 | 20 | 6.0 |
| 5-Jul | 4 | 15 | 10.7 | 2 | south | flood | 30.7 | 24 | 7.0 |
| | 5 | 13 | 10.4 | 2 | south | flood | 31.2 | 37 | 8.0 |
| | 6 | 13 | 10.3 | 2 | south | flood | 31.1 | 48 | 5.0 |
| | 6.5 | 15 | 12.8 | 2 | south | flood | 27.9 | 44 | 3.0 |
| | 7 | 14 | 13.0 | 2 | south | ebb | 27.9 | 40 | 4.5 |
| | 8 | 12 | 11.6 | 2 | south | ebb | 29.0 | 30 | 5.0 |
| 6-Jul | 8 | 11 | 11.5 | 5 | south | ebb | 29.8 | 29 | 3.0 |
| | 7 | 12 | 11.5 | 5 | south | ebb | 28.6 | 44 | 2.5 |
| | 6.5 | 14 | 11.4 | 5 | south | ebb | 28.7 | 41 | 4.5 |
| | 6 | 14 | 10.8 | 2 | south | ebb | 29.8 | 44 | 4.5 |
| | 5 | 17 | 10.4 | 2 | south | flood | 30.9 | 36 | 5.5 |
| | 4 | 13 | 10.4 | 5 | south | flood | 31.3 | 26 | 7.0 |

-continued-

Appendix A14.–Page 2 of 5.

| Date | Sta | Air Temp (c) | Water Temp (c) | Wind Vel. (knots) | Wind Dir | Tide Stage | Salinity (ppt) | Water Depth (f) | Secchi (m) |
|--------|----------------|--------------|----------------|-------------------|-----------|------------|----------------|-----------------|------------|
| 7-Jul | 4 | 12 | 10.5 | 4 | south | low | 31.1 | 24 | 5.5 |
| | 5 | 13 | 10.5 | 4 | south | flood | 30.9 | 36 | 6.0 |
| | 6 | 13 | 10.8 | 4 | south | flood | 29.7 | 46 | 3.5 |
| | 6.5 | 12 | 12.4 | 4 | south | flood | 29.5 | 44 | 3.5 |
| | 7 | 13 | 11.6 | 8 | south | flood | 28.8 | 46 | 2.5 |
| | 8 | 13 | 11.5 | 8 | south | ebb | 29.2 | 29 | 2.0 |
| 8-Jul | 8 | 11 | 11.4 | 10 | southwest | high | 29.0 | 32 | 2.5 |
| | 7 | 11 | 11.4 | 15 | southwest | ebb | 29.0 | 45 | 2.5 |
| | 6.5 | 12 | 11.6 | 15 | southwest | ebb | 28.8 | 44 | 2.5 |
| | 6 | 13 | 11.2 | 15 | southwest | ebb | 29.2 | 46 | 2.0 |
| | 5 | 14 | 12.1 | 10 | south | ebb | 28.6 | 33 | 5.0 |
| | 4 | 12 | 10.8 | 5 | south | ebb | 30.9 | 24 | 6.0 |
| 9-Jul | 4 | 14 | 10.6 | 1 | south | ebb | 31.4 | 24 | 7.5 |
| | 5 | 17 | 13.3 | 1 | south | ebb | 28.4 | 35 | 4.5 |
| | 6 | 16 | 11.7 | 1 | south | ebb | 29.0 | 46 | 4.0 |
| | 6.5 | 15 | 12.6 | 3 | south | flood | 30.6 | 45 | 4.5 |
| | 7 | 15 | 13.0 | 9 | south | flood | 27.9 | 49 | 4.0 |
| | 8 | 15 | 12.0 | 4 | south | flood | 28.0 | 31 | 2.0 |
| 10-Jul | 8 | 12 | 11.6 | 25 | southwest | flood | 28.9 | 31 | 2.0 |
| | 7 | 12 | 11.6 | 15 | southwest | ebb | 28.9 | 45 | 4.0 |
| | 6.5 | 12 | 11.6 | 15 | southwest | ebb | 29.0 | 44 | 3.0 |
| | 6 | 12 | 11.6 | 20 | southeast | ebb | 29.2 | 48 | 4.0 |
| | 5 ^a | - | - | - | - | - | - | - | - |
| | 4 ^a | - | - | - | - | - | - | - | - |
| 11-Jul | 4 | 13 | 10.9 | 7 | southwest | ebb | 30.3 | 24 | 3.5 |
| | 5 | 14 | 11.6 | 9 | west | ebb | 29.3 | 35 | 4.0 |
| | 6 | 13 | 11.6 | 9 | south | ebb | 29.3 | 47 | 4.0 |
| | 6.5 | 12 | 12.7 | 9 | south | ebb | 28.3 | 41 | 3.5 |
| | 7 | 12 | 12.4 | 9 | south | flood | 28.2 | 45 | 3.0 |
| | 8 | 12 | 12.2 | 10 | south | flood | 28.8 | 29 | 4.0 |
| 12-Jul | 8 | 11 | 11.6 | 2 | southeast | high | 29.1 | 32 | 3.5 |
| | 7 | 11 | 11.6 | 2 | southeast | high | 29.3 | 46 | 2.5 |
| | 6.5 | 12 | 12.9 | 8 | southeast | ebb | 28.8 | 46 | 3.5 |
| | 6 | 12 | 11.8 | 8 | southeast | ebb | 29.1 | 45 | 4.0 |
| | 5 | 12 | 12.1 | 2 | southeast | flood | 28.6 | 35 | 4.0 |
| | 4 | 12 | 12.3 | 2 | southeast | ebb | 28.7 | 24 | 4.0 |

-continued-

Appendix A14.–Page 3 of 5.

| Date | Sta | Air Temp (c) | Water Temp (c) | Wind Vel. (knots) | Wind Dir | Tide Stage | Salinity (ppt) | Water Depth (f) | Secchi (m) |
|--------|-----|--------------|----------------|-------------------|-----------|------------|----------------|-----------------|------------|
| 13-Jul | 4 | 11 | 10.2 | 11 | northeast | ebb | 31.5 | 25 | 8.5 |
| | 5 | 13 | 11.6 | 7 | northeast | ebb | 29.7 | 37 | 4.0 |
| | 6 | 12 | 11.9 | 12 | north | ebb | 29.5 | 44 | 3.5 |
| | 6.5 | 13 | 11.6 | 8 | northeast | ebb | 27.9 | 43 | 4.5 |
| | 7 | 14 | 12.1 | 1 | northwest | low | 29.3 | 43 | 3.5 |
| | 8 | 13 | 12.3 | 8 | northwest | low | 28.8 | 26 | 3.5 |
| 14-Jul | 8 | 13 | 12.4 | 8 | northwest | low | 28.0 | 30 | 3.0 |
| | 7 | 13 | 11.3 | 8 | northwest | flood | 29.9 | 42 | 5.2 |
| | 6.5 | 12 | 11.1 | 6 | northwest | flood | 30.5 | 45 | 6.0 |
| | 6 | 14 | 11.5 | 6 | northwest | flood | 30.1 | 46 | 7.0 |
| | 5 | 18 | 10.1 | 1 | east | ebb | 31.5 | 38 | 7.5 |
| | 4 | 19 | 10.4 | 3 | northwest | high | 31.5 | 24 | 10.0 |
| 15-Jul | 4 | 12 | 10.6 | 9 | southwest | ebb | 31.6 | 25 | 10.0 |
| | 5 | 12 | 11.1 | 8 | southwest | flood | 31.6 | 37 | 6.0 |
| | 6 | 14 | 13.5 | 12 | south | flood | 26.5 | 48 | 2.5 |
| | 6.5 | 17 | 13.3 | 9 | south | flood | 29.5 | 43 | 2.5 |
| | 7 | 15 | 13.8 | 12 | south | ebb | 26.2 | 45 | 4.5 |
| | 8 | 15 | 13.5 | 13 | south | ebb | 26.7 | 26 | 4.0 |
| 16-Jul | 8 | 13 | 13.1 | 12 | northeast | flood | 26.1 | 29 | 3.5 |
| | 7 | 13 | 13.8 | 13 | northeast | flood | 25.5 | 46 | 3.5 |
| | 6.5 | 13 | 12.7 | 15 | north | flood | 27.8 | 44 | 4.0 |
| | 6 | 12 | 10.1 | 15 | north | high | 31.4 | 49 | 7.0 |
| | 5 | 13 | 10.3 | 12 | north | high | 31.4 | 37 | 7.0 |
| | 4 | 12 | 10.3 | 11 | northeast | high | 31.5 | 24 | 9.5 |
| 17-Jul | 4 | 15 | 10.4 | 4 | southwest | flood | 31.5 | 25 | 9.0 |
| | 5 | 13 | 10.3 | 5 | south | flood | 31.5 | 37 | 8.5 |
| | 6 | 13 | 12.4 | 7 | southwest | flood | 28.2 | 48 | 6.5 |
| | 6.5 | 13 | 13.5 | 11 | southwest | flood | 25.7 | 44 | 4.5 |
| | 7 | 14 | 14.5 | 8 | southwest | high | 26.2 | 46 | 4.5 |
| | 8 | 14 | 12.8 | 7 | south | high | 27.3 | 27 | 4.5 |
| 18-Jul | 8 | 14 | 13.2 | 6 | south | ebb | 26.6 | 29 | 4.5 |
| | 7 | 15 | 13.5 | 5 | south | ebb | 26.6 | 42 | 4.5 |
| | 6.5 | 15 | 12.6 | 4 | south | ebb | 27.6 | 43 | 7.0 |
| | 6 | 12 | 11.1 | 5 | south | low | 30.5 | 46 | 8.5 |
| | 5 | 12 | 10.6 | 4 | southeast | low | 31.1 | 48 | 8.5 |
| | 4 | 16 | 10.5 | 2 | south | low | 31.3 | 36 | 10.0 |

-continued-

Appendix A14.–Page 4 of 5.

| Date | Sta | Air | Water | Wind | Wind | Tide | Salinity | Water | Secchi | |
|--------|--------|------|-------|---------|-----------|-----------|----------|-------|--------|-------|
| | | Temp | Temp | Vel. | | | | Dir | | Depth |
| | | (c) | (c) | (knots) | | | (ppt) | (f) | (m) | |
| 19-Jul | 4 | 14 | 10.7 | 1 | northwest | flood | 31.4 | 25 | 7.0 | |
| | 5 | 13 | 10.3 | 2 | southwest | flood | 31.5 | 38 | 8.0 | |
| | 6 | 13 | 12.6 | 3 | southwest | flood | 28.2 | 25 | 7.0 | |
| | 6.5 | 14 | 12.6 | 3 | southwest | high | 27.8 | 42 | 6.0 | |
| | 7 | 13 | 13.1 | 3 | southwest | high | 26.6 | 45 | 3.5 | |
| | 8 | 14 | 12.7 | 4 | southwest | ebb | 27.3 | 30 | 4.5 | |
| | 20-Jul | 8 | 14 | 11.9 | 4 | south | ebb | 28.3 | 29 | 4.5 |
| | | 7 | 15 | 12.9 | 2 | west | ebb | 26.7 | 42 | 3.5 |
| 6.5 | | 12 | 11.5 | 4 | southeast | low | 29.5 | 42 | 6.0 | |
| 6 | | 15 | 10.6 | 3 | southeast | flood | 31.1 | 47 | 10.0 | |
| 5 | | 14 | 10.4 | 1 | southwest | flood | 31.5 | 38 | 9.0 | |
| 21-Jul | 4 | 14 | 10.4 | 4 | southwest | flood | 31.6 | 26 | 12.5 | |
| | 4 | 14 | 10.8 | 14 | south | ebb | 31.3 | 22 | 6.5 | |
| | 5 | 12 | 10.6 | 11 | south | flood | 31.3 | 35 | 5.5 | |
| | 6 | 12 | 10.7 | 6 | south | flood | 30.6 | 45 | 5.5 | |
| | 6.5 | 13 | 11.0 | 3 | south | flood | 30.3 | 42 | 5.0 | |
| | 7 | 13 | 11.5 | 4 | south | flood | 29.7 | 46 | 5.5 | |
| | 8 | 13 | 11.7 | 6 | south | flood | 29.4 | 32 | 3.5 | |
| | 22-Jul | 8 | 13 | 11.7 | 4 | southwest | high | 28.8 | 33 | 2.5 |
| 7 | | 14 | 12.0 | 8 | southwest | ebb | 28.5 | 46 | 2.5 | |
| 6.5 | | 18 | 12.0 | 1 | south | ebb | 28.8 | 42 | 3.5 | |
| 6 | | 15 | 11.7 | 3 | southeast | ebb | 29.9 | 44 | 5.0 | |
| 5 | | 18 | 11.4 | 1 | south | low | 31.1 | 35 | 9.0 | |
| 4 | | 11 | 12.7 | 4 | southeast | flood | 31.2 | 23 | 7.2 | |
| 23-Jul | | 4 | 13 | 10.8 | 2 | southwest | ebb | 31.5 | 22 | 7.0 |
| | 5 | 13 | 11.0 | 2 | southwest | ebb | 31.1 | 34 | 7.0 | |
| | 6 | 13 | 11.5 | 4 | southwest | low | 32.0 | 46 | 2.0 | |
| | 6.5 | 15 | 11.5 | 4 | southwest | low | 30.1 | 43 | 3.0 | |
| | 7 | 13 | 11.7 | 6 | southwest | flood | 30.2 | 46 | 4.5 | |
| | 8 | 13 | 11.2 | 5 | southwest | flood | 30.2 | 33 | 3.0 | |
| | 24-Jul | 8 | 14 | 11.2 | 5 | north | flood | 30.0 | 33 | 3.0 |
| 7 | | 12 | 11.2 | 4 | southeast | high | 30.2 | 43 | 4.0 | |
| 6.5 | | 13 | 11.2 | 5 | north | ebb | 30.3 | 42 | 4.0 | |
| 6 | | 13 | 11.5 | 7 | north | ebb | 30.0 | 44 | 3.0 | |
| 5 | | 12 | 11.5 | 9 | north | ebb | 31.1 | 34 | 5.0 | |
| 4 | | 12 | 11.1 | 16 | north | low | 31.2 | 23 | 6.0 | |

-continued-

Appendix A14.–Page 5 of 5.

| Date | Sta | Air | Water | Wind | Wind Dir | Tide Stage | Salinity (ppt) | Water | Secchi (m) | |
|--------|--------|----------|----------|--------------|-----------|------------|----------------|-----------|------------|-----|
| | | Temp (c) | Temp (c) | Vel. (knots) | | | | Depth (f) | | |
| 25-Jul | 4 | 12 | 10.5 | 14 | northwest | ebb | 31.4 | 23 | 6.5 | |
| | 5 | 12 | 11.2 | 16 | northwest | ebb | 30.8 | 35 | 5.0 | |
| | 6 | 12 | 11.6 | 16 | northwest | ebb | 30.2 | 47 | 4.5 | |
| | 6.5 | 14 | 11.8 | 2 | northeast | flood | 29.9 | 43 | 5.4 | |
| | 7 | 12 | 12.0 | 3 | east | flood | 29.7 | 46 | 2.5 | |
| | 8 | 14 | 12.0 | 1 | south | flood | 29.6 | 28 | 3.0 | |
| | 26-Jul | 8 | 14 | 12.0 | 6 | northwest | flood | 29.1 | 30 | 2.0 |
| | | 7 | 12 | 11.5 | 6 | north | flood | 30.2 | 47 | 3.5 |
| 6.5 | | 12 | 11.7 | 7 | north | high | 30.8 | 44 | 5.5 | |
| 6 | | 11 | 10.9 | 7 | east | ebb | 31.1 | 49 | 7.5 | |
| 5 | | 12 | 12.0 | 4 | north | ebb | 31.5 | 35 | 4.5 | |
| 4 | | 15 | 11.1 | 4 | northwest | ebb | 30.2 | 23 | 6.5 | |
| 27-Jul | 4 | 10 | 12.6 | 4 | north | ebb | 31.4 | 24 | 9.0 | |
| | 5 | 12 | 11.4 | 12 | northwest | ebb | 30.6 | 36 | 5.0 | |
| | 6 | 13 | 12.9 | 12 | west | low | 28.9 | 48 | 2.5 | |
| | 6.5 | 13 | 12.9 | 15 | northwest | ebb | 28.6 | 43 | 3.5 | |
| | 7 | 14 | 12.9 | 14 | west | flood | 28.7 | 46 | 3.5 | |
| | 8 | 12 | 14.4 | 13 | northwest | flood | 29.0 | 28 | 2.5 | |
| 28-Jul | 8 | 13 | 12.8 | 5 | southwest | low | 28.2 | 31 | 3.0 | |
| | 7 | 12 | 12.5 | 8 | southeast | flood | 29.0 | 45 | 3.0 | |
| | 6.5 | 12 | 11.6 | 9 | southeast | flood | 30.2 | 44 | 5.0 | |
| | 6 | 13 | 10.8 | 8 | southeast | flood | 30.1 | 48 | 8.0 | |
| | 5 | 11 | 10.5 | 6 | east | flood | 32.3 | 37 | 11.0 | |
| | 4 | 15 | 12.3 | 10 | east | ebb | 30.3 | 25 | 12.0 | |
| 29-Jul | 4 | 11 | 10.3 | 2 | south | flood | 31.4 | 26 | 9.5 | |
| | 5 | 11 | 10.4 | 2 | south | flood | 31.0 | 37 | 11.5 | |
| | 6 | 12 | 13.5 | 6 | south | high | 27.0 | 43 | 3.5 | |
| | 6.5 | 13 | 13.5 | 8 | southwest | ebb | 27.1 | 43 | 4.0 | |
| | 7 | 14 | 13.5 | 9 | southeast | ebb | 27.4 | 45 | 3.5 | |
| | 8 | 13 | 13.0 | 9 | southwest | ebb | 26.3 | 28 | 2.8 | |
| 30-Jul | 8 | 13 | 12.9 | 15 | southwest | ebb | 28.2 | 30 | 2.8 | |
| | 7 | 13 | 13.2 | 17 | southwest | ebb | 26.9 | 45 | 3.5 | |
| | 6.5 | 12 | 13.2 | 17 | southeast | ebb | 26.8 | 43 | 3.0 | |
| | 6 | 11 | 12.9 | 19 | southeast | ebb | 28.5 | 50 | 3.0 | |
| | 5 | 12 | 13.4 | 18 | south | flood | 27.6 | 35 | 4.0 | |
| | 4 | 13 | 12.8 | 20 | southeast | flood | 28.8 | 24 | 4.5 | |

Appendix A15.—Yearly mean values of physical observations made during the 1995–2005 offshore test fish project.

| Sta | Year | Air | Water | Wind | Wind | Water | | | Sta | Year | Air | Water | Wind | Wind | Water | | |
|-----|------|------|-------|---------|------|-------|----------|-------|-----|------|--------|-------|---------|------|-------|------|----------|
| | | Temp | Temp | Vel. | | Dir | Salinity | Depth | | | Secchi | Temp | Temp | | Vel. | Dir | Salinity |
| | | (c) | (c) | (knots) | | (ppt) | (f) | (m) | | | (c) | (c) | (knots) | | (ppt) | (f) | (m) |
| 4 | 1996 | 12.8 | 9.4 | 7.4 | E | 31.5 | 24.6 | 7.9 | 6.5 | 1996 | 13.2 | 10.3 | 8.4 | E | 30.5 | 42.9 | 3.2 |
| | 1997 | 13.7 | 9.5 | 8.9 | SE | 31.6 | 24.6 | 6.2 | | 1997 | 13.9 | 10.7 | 9.4 | E | 29.6 | 43.4 | 3.2 |
| | 1998 | 12.5 | 9.7 | 9.7 | SE | 31.0 | 24.4 | 9.5 | | 1998 | 12.7 | 10.5 | 7.7 | SE | 29.5 | 43.3 | 3.5 |
| | 1999 | 13.1 | 9.6 | 10.6 | SE | 31.4 | 24.3 | 7.6 | | 1999 | 13.4 | 10.5 | 13.0 | SE | 29.7 | 43.2 | 3.5 |
| | 2000 | 13.8 | 9.7 | 10.0 | SE | 31.5 | 23.5 | 10.0 | | 2000 | 13.6 | 10.8 | 13.0 | S | 29.7 | 42.9 | 3.7 |
| | 2001 | 12.9 | 9.8 | 11.1 | SE | 31.5 | 23.6 | 8.4 | | 2001 | 12.8 | 11.1 | 11.8 | S | 29.4 | 42.7 | 4.0 |
| | 2002 | 12.6 | 9.5 | 12.6 | S | 31.4 | 23.6 | 8.1 | | 2002 | 12.6 | 10.4 | 13.7 | S | 30.0 | 42.6 | 3.3 |
| | 2003 | 14.1 | 10.6 | 12.0 | S | 31.2 | 23.4 | 8.3 | | 2003 | 14.4 | 11.7 | 14.9 | S | 29.1 | 41.3 | 4.1 |
| | 2004 | 10.7 | 9.6 | 7.1 | E | 31.3 | 23.8 | 7.9 | | 2004 | 10.7 | 10.8 | 10.1 | SE | 29.4 | 41.6 | 3.6 |
| | 2005 | 12.9 | 10.9 | 6.2 | S | 31.0 | 24.5 | 7.4 | | 2005 | 13.2 | 12.2 | 7.4 | S | 28.7 | 42.8 | 4.2 |
| Avg | | 12.9 | 9.8 | 9.6 | SE | 31.4 | 24.0 | 8.1 | Avg | | 13.1 | 10.9 | 10.9 | SE | 29.6 | 42.7 | 3.6 |
| 5 | 1996 | 12.8 | 9.4 | 8.6 | NE | 31.3 | 36.3 | 6.3 | 7 | 1996 | 12.7 | 10.4 | 10.7 | SE | 30.3 | 44.9 | 3.4 |
| | 1997 | 13.7 | 9.9 | 10.1 | SE | 31.3 | 36.8 | 5.2 | | 1997 | 14.0 | 10.9 | 10.3 | SE | 30.2 | 44.8 | 2.9 |
| | 1998 | 12.8 | 9.8 | 9.8 | SE | 31.1 | 35.2 | 8.5 | | 1998 | 12.3 | 10.7 | 8.4 | SE | 29.1 | 44.3 | 3.0 |
| | 1999 | 13.4 | 10.0 | 12.9 | SE | 30.6 | 38.9 | 6.2 | | 1999 | 13.3 | 10.6 | 13.0 | S | 29.5 | 42.7 | 2.9 |
| | 2000 | 13.5 | 10.1 | 11.8 | SE | 30.7 | 35.9 | 7.1 | | 2000 | 13.1 | 10.9 | 13.6 | S | 29.4 | 43.3 | 3.0 |
| | 2001 | 12.9 | 10.1 | 11.2 | SE | 31.0 | 35.5 | 6.9 | | 2001 | 13.1 | 11.4 | 9.9 | SE | 29.0 | 43.6 | 3.5 |
| | 2002 | 12.8 | 9.7 | 13.9 | S | 30.9 | 35.8 | 6.3 | | 2002 | 12.4 | 10.4 | 12.4 | SE | 29.9 | 44.0 | 2.8 |
| | 2003 | 14.0 | 11.0 | 13.3 | SE | 30.6 | 35.7 | 6.3 | | 2003 | 14.3 | 11.6 | 13.0 | S | 29.0 | 44.3 | 3.6 |
| | 2004 | 10.7 | 9.9 | 7.2 | SE | 30.7 | 34.7 | 7.1 | | 2004 | 10.6 | 11.0 | 9.7 | SE | 28.8 | 44.7 | 2.7 |
| | 2005 | 13.1 | 11.1 | 5.9 | S | 30.6 | 36.3 | 6.5 | | 2005 | 12.9 | 12.3 | 7.6 | S | 28.3 | 44.8 | 3.6 |
| Avg | | 13.0 | 10.1 | 10.4 | SE | 30.9 | 36.1 | 6.6 | Avg | | 12.9 | 11.0 | 10.8 | SE | 29.3 | 44.1 | 3.1 |

-continued-

Appendix A15.–Page 2 of 2.

| Sta | Year | Air | Water | Wind | | Water | | |
|-----|------|------|-------|---------|------|----------|-------|--------|
| | | Temp | Temp | Vel. | Wind | Salinity | Depth | Secchi |
| | | (c) | (c) | (knots) | Dir | (ppt) | (f) | (m) |
| 6 | 1996 | 12.4 | 10.2 | 9.7 | E | 30.6 | 47.1 | 4.2 |
| | 1997 | 13.8 | 10.5 | 11.1 | SE | 30.8 | 45.9 | 3.7 |
| | 1998 | 12.4 | 10.3 | 10.9 | S | 30.0 | 46.1 | 4.7 |
| | 1999 | 13.5 | 10.3 | 12.5 | SE | 29.8 | 44.4 | 4.3 |
| | 2000 | 13.5 | 10.6 | 11.1 | SE | 29.9 | 45.4 | 4.9 |
| | 2001 | 12.8 | 10.7 | 10.7 | S | 30.5 | 46.2 | 5.2 |
| | 2002 | 12.8 | 10.1 | 13.4 | S | 30.4 | 45.1 | 4.2 |
| | 2003 | 14.7 | 11.5 | 12.9 | S | 29.5 | 46.4 | 4.9 |
| | 2004 | 10.6 | 10.3 | 8.0 | SE | 30.1 | 46.6 | 4.6 |
| | 2005 | 12.8 | 11.6 | 8.0 | S | 29.4 | 45.8 | 4.7 |
| Avg | | 12.9 | 10.6 | 10.8 | SE | 30.1 | 45.9 | 4.5 |

| Sta | Year | Air | Water | Wind | | Water | | |
|-----|------|------|-------|---------|------|----------|-------|--------|
| | | Temp | Temp | Vel. | Wind | Salinity | Depth | Secchi |
| | | (c) | (c) | (knots) | Dir | (ppt) | (f) | (m) |
| 8 | 1996 | 12.2 | 10.4 | 9.3 | SE | 30.3 | 29.9 | 2.7 |
| | 1997 | 13.7 | 11.1 | 9.6 | SE | 30.1 | 30.1 | 2.6 |
| | 1998 | 12.5 | 10.7 | 9.1 | S | 29.1 | 29.3 | 2.8 |
| | 1999 | 13.6 | 10.5 | 11.8 | SE | 30.0 | 25.9 | 2.6 |
| | 2000 | 13.2 | 11.0 | 14.0 | S | 29.5 | 29.1 | 2.6 |
| | 2001 | 12.8 | 11.3 | 9.5 | SE | 29.0 | 28.9 | 3.1 |
| | 2002 | 12.1 | 10.3 | 11.8 | SE | 30.0 | 29.4 | 2.4 |
| | 2003 | 13.7 | 11.2 | 11.6 | SE | 28.1 | 28.9 | 3.1 |
| | 2004 | 10.8 | 11.0 | 9.1 | SE | 29.3 | 28.7 | 2.4 |
| | 2005 | 12.8 | 12.1 | 7.7 | S | 28.5 | 29.8 | 3.3 |
| Avg | | 12.8 | 11.0 | 10.3 | S | 29.4 | 29.0 | 2.8 |

Appendix A16.—Yearly mean values for selected chemical and physical variables collected during the offshore test fish project, 1979–2005.

| Year | Air Temp. (c) | Water Temp. (c) | Wind Vel. (knots) | Salinity (ppt) | Secchi (m) |
|---------------|------------------------------|--------------------------------|----------------------------------|---------------------------|-----------------------|
| 1979 | 12.4 | 12.2 | 5.9 | 25.0 | 5.7 |
| 1980 | 12.4 | 10.0 | 8.2 | 24.8 | 4.2 |
| 1981 | 13.4 | 11.0 | 10.1 | 23.1 | 4.1 |
| 1982 | 12.0 | 8.5 | 9.0 | 20.3 | 5.0 |
| 1983 | 14.9 | 10.9 | 9.4 | 20.6 | 4.7 |
| 1984 | 13.5 | 10.8 | 9.1 | 14.3 | 5.3 |
| 1985 | 10.8 | 8.2 | 9.2 | 28.0 | 5.5 |
| 1986 | 10.6 | 9.1 | 8.2 | - | 5.4 |
| 1987 | 12.6 | 10.1 | 4.1 | 28.4 | 5.1 |
| 1988 | 14.2 | 9.1 | 8.9 | 30.2 | 4.7 |
| 1989 | 13.1 | 10.0 | 4.4 | 27.7 | 4.7 |
| 1990 | 12.3 | 11.4 | 8.5 | 21.3 | 4.6 |
| 1991 | 10.9 | 9.9 | 6.6 | 13.1 | 4.1 |
| 1992 | 12.0 | 11.1 | 5.4 | 28.4 | 4.3 |
| 1993 | 13.5 | 10.5 | 6.9 | 26.2 | 5.0 |
| 1994 | 13.0 | 10.0 | 9.3 | 29.0 | 6.0 |
| 1995 | 13.1 | 9.5 | 7.9 | 26.5 | 4.6 |
| 1996 | 12.6 | 10.0 | 9.1 | 30.8 | 4.7 |
| 1997 | 13.8 | 10.5 | 10.0 | 30.6 | 4.0 |
| 1998 | 12.5 | 10.3 | 8.3 | 30.0 | 5.4 |
| 1999 | 13.4 | 10.3 | 12.4 | 30.2 | 4.5 |
| 2000 | 13.5 | 10.5 | 12.2 | 30.1 | 5.2 |
| 2001 | 12.9 | 10.7 | 10.7 | 30.1 | 5.2 |
| 2002 | 12.5 | 10.1 | 13.0 | 30.4 | 4.5 |
| 2003 | 14.2 | 11.3 | 12.9 | 29.6 | 5.0 |
| 2004 | 10.7 | 10.4 | 8.5 | 30.0 | 4.7 |
| 1992–2004 Avg | 12.9 | 10.4 | 9.7 | 29.4 | 4.9 |
| 2005 | 13.0 | 11.7 | 7.1 | 29.4 | 5.0 |